

=> FILE HCAPLUS

FILE 'HCAPLUS' ENTERED AT 12:08:01 ON 27 MAY 2003

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2003 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 27 May 2003 VOL 138 ISS 22

FILE LAST UPDATED: 26 May 2003 (20030526/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> D QUE L30

L2	1457740	SEA FILE=HCAPLUS ABB=ON	RAT OR RATS OR RATTUS OR MOUSE OR MICE OR RODENT#
L3	34259	SEA FILE=HCAPLUS ABB=ON	L2 AND (TEST? OR MEASUR? OR SCREEN?) AND (RODENTIC? OR TOXIC?)
L4	1	SEA FILE=REGISTRY ABB=ON	9004-34-6
L5	394825	SEA FILE=HCAPLUS ABB=ON	L4 OR CORNCOB? OR COB OR COBS OR MAIZE OR ?CELLULOS?
L7	138348	SEA FILE=HCAPLUS ABB=ON	L2 AND (FEED? OR DIET?)
L8	2130	SEA FILE=HCAPLUS ABB=ON	L5 AND L7
L9	47045	SEA FILE=HCAPLUS ABB=ON	(WEIGHT? OR WT) (2A) LOSS
L10	59149	SEA FILE=HCAPLUS ABB=ON	L9 OR AD LIBITUM OR FREE? (2A) FEED?
L11	47	SEA FILE=HCAPLUS ABB=ON	L8 AND L3
L12	3	SEA FILE=HCAPLUS ABB=ON	L10 AND L11
L13	246	SEA FILE=HCAPLUS ABB=ON	L3 AND L5
L14	3	SEA FILE=HCAPLUS ABB=ON	L10 AND L13
L15	666145	SEA FILE=HCAPLUS ABB=ON	BLOAT? OR IMPACT? OR COMPACT? OR CAECUM OR INTESTIN? OR GUT OR STOMACH? OR CECUM
L16	2445	SEA FILE=HCAPLUS ABB=ON	L3 AND L15
L18	474846	SEA FILE=HCAPLUS ABB=ON	L5 OR CORN
L19	347	SEA FILE=HCAPLUS ABB=ON	L7 AND L18 AND L10
L20	71	SEA FILE=HCAPLUS ABB=ON	L15 AND L19
L21	0	SEA FILE=HCAPLUS ABB=ON	L20 AND (WATER? (2A) (RETEN? OR RETAIN?) OR ION(W) TRANSPORT? OR POST MORTEM OR AUTOPS? OR NECROPS?)
L22	65	SEA FILE=HCAPLUS ABB=ON	L16 AND L18
L23	2	SEA FILE=HCAPLUS ABB=ON	L20 AND L3
L24	4	SEA FILE=HCAPLUS ABB=ON	L22 AND (WATER? (2A) (RETEN? OR RETAIN?) OR ION(W) TRANSPORT? OR POST MORTEM OR AUTOPS? OR NECROPS?)
L25	9	SEA FILE=HCAPLUS ABB=ON	L12 OR L14 OR L21 OR L23 OR L24
L26	12	SEA FILE=HCAPLUS ABB=ON	L3 AND L7 AND L10 AND L18
L27	65	SEA FILE=HCAPLUS ABB=ON	L3 AND L18 AND L15
L28	7	SEA FILE=HCAPLUS ABB=ON	L27 AND L10

L29 4 SEA FILE=HCAPLUS ABB=ON L27 AND (WATER?(2A) (RETEN? OR
RETAIN?) OR ION(W)TRANSPORT? OR POST MORTEM OR AUTOPS? OR
NECROPS?)
L30 20 SEA FILE=HCAPLUS ABB=ON L25 OR L26 OR L28 OR L29

=> FILE WPIX

FILE 'WPIX' ENTERED AT 12:08:09 ON 27 MAY 2003
COPYRIGHT (C) 2003 THOMSON DERWENT

FILE LAST UPDATED: 26 MAY 2003 <20030526/UP>
MOST RECENT DERWENT UPDATE: 200333 <200333/DW>
DERWENT WORLD PATENTS INDEX SUBSCRIBER FILE, COVERS 1963 TO DATE

>>> NEW WEEKLY SDI FREQUENCY AVAILABLE --> see NEWS <<<

>>> SLART (Simultaneous Left and Right Truncation) is now
available in the /ABEX field. An additional search field
/BIX is also provided which comprises both /BI and /ABEX <<<

>>> PATENT IMAGES AVAILABLE FOR PRINT AND DISPLAY <<<

>>> FOR DETAILS OF THE PATENTS COVERED IN CURRENT UPDATES,
SEE <http://www.derwent.com/dwpi/updates/dwpicov/index.html> <<<

>>> FOR A COPY OF THE DERWENT WORLD PATENTS INDEX STN USER GUIDE,
PLEASE VISIT:
http://www.stn-international.de/training_center/patents/stn_guide.pdf <<<

>>> FOR INFORMATION ON ALL DERWENT WORLD PATENTS INDEX USER
GUIDES, PLEASE VISIT:
http://www.derwent.com/userguides/dwpi_guide.html <<<

=> D QUE L38

L2 1457740 SEA FILE=HCAPLUS ABB=ON RAT OR RATS OR RATTUS OR MOUSE OR
MICE OR RODENT#
L4 1 SEA FILE=REGISTRY ABB=ON 9004-34-6
L5 394825 SEA FILE=HCAPLUS ABB=ON L4 OR CORNCOB? OR COB OR COBS OR
MAIZE OR ?CELLULOS?
L31 1658 SEA FILE=WPIX ABB=ON L2 AND (TEST? OR MEASUR? OR SCREEN?) AND
(RODENTIC? OR TOXIC?)
L32 131261 SEA FILE=WPIX ABB=ON L5 OR CORN
L33 63 SEA FILE=WPIX ABB=ON L31 AND L32
L34 339776 SEA FILE=WPIX ABB=ON BLOAT? OR IMPACT? OR COMPACT? OR CAECUM
OR INTESTIN? OR GUT OR STOMACH? OR CECUM
L35 11 SEA FILE=WPIX ABB=ON L33 AND L34
L36 2 SEA FILE=WPIX ABB=ON L33 AND (WATER?(2A) (RETEN? OR RETAIN?)
OR ION(W)TRANSPORT? OR POST MORTEM OR AUTOPS? OR NECROPS?)
L37 12 SEA FILE=WPIX ABB=ON L35 OR L36
L38 2 SEA FILE=WPIX ABB=ON L37 AND A01N?/IC

=> FILE CABA

FILE 'CABA' ENTERED AT 12:08:20 ON 27 MAY 2003
COPYRIGHT (C) 2003 CAB INTERNATIONAL (CABI)

FILE COVERS 1973 TO 2 May 2003 (20030502/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> D QUE L52

L2 1457740 SEA FILE=HCAPLUS ABB=ON RAT OR RATS OR RATTUS OR MOUSE OR
MICE OR RODENT#
L4 1 SEA FILE=REGISTRY ABB=ON 9004-34-6
L5 394825 SEA FILE=HCAPLUS ABB=ON L4 OR CORNCOB? OR COB OR COBS OR
MAIZE OR ?CELLULOS?
L31 1658 SEA FILE=WPIX ABB=ON L2 AND (TEST? OR MEASUR? OR SCREEN?) AND
(RODENTIC? OR TOXIC?)
L32 131261 SEA FILE=WPIX ABB=ON L5 OR CORN
L33 63 SEA FILE=WPIX ABB=ON L31 AND L32
L34 339776 SEA FILE=WPIX ABB=ON BLOAT? OR IMPACT? OR COMPACT? OR CAECUM
OR INTESTIN? OR GUT OR STOMACH? OR CECUM
L35 11 SEA FILE=WPIX ABB=ON L33 AND L34
L36 2 SEA FILE=WPIX ABB=ON L33 AND (WATER?(2A) (RETEN? OR RETAIN?)
OR ION(W)TRANSPORT? OR POST MORTEM OR AUTOPS? OR NECROPS?)
L39 30 SEA FILE=CABA ABB=ON L35 OR L36
L40 0 SEA FILE=CABA ABB=ON L39 AND RODENTICID?
L41 232245 SEA FILE=CABA ABB=ON RAT OR RATS OR RATTUS OR MOUSE OR MICE
OR RODENT#
L43 2712 SEA FILE=CABA ABB=ON RODENTICIDES+NT/CT
L44 380 SEA FILE=CABA ABB=ON RODENT CONTROL+NT/CT
L45 1181 SEA FILE=CABA ABB=ON L41 AND (L43 OR L44)
L46 219115 SEA FILE=CABA ABB=ON BLOAT? OR IMPACT? OR COMPACT? OR CAECUM
OR INTESTIN? OR GUT OR STOMACH? OR CECUM
L47 65 SEA FILE=CABA ABB=ON L45 AND L46
L48 0 SEA FILE=CABA ABB=ON L47 AND CELLULOS?
L49 0 SEA FILE=CABA ABB=ON L48 AND (WATER?(2A) (RETEN? OR RETAIN?)
OR ION(W)TRANSPORT? OR POST MORTEM OR AUTOPS? OR NECROPS?)
L50 0 SEA FILE=CABA ABB=ON L39 AND (L43 OR L44)
L51 0 SEA FILE=CABA ABB=ON L47 AND NONTOXIC?
L52 0 SEA FILE=CABA ABB=ON L40 OR L48 OR L49 OR L50 OR L51

=> FILE AGRICOLA

FILE 'AGRICOLA' ENTERED AT 12:08:33 ON 27 MAY 2003

FILE COVERS 1970 TO 15 May 2003 (20030515/ED)

Compiled and distributed by the National Agricultural Library
of the Department of Agriculture of the United States of
America. It contains copyrighted material. All rights
reserved. (2003)

This file contains CAS Registry Numbers for easy and accurate
substance identification.

=> D QUE L62

L2 1457740 SEA FILE=HCAPLUS ABB=ON RAT OR RATS OR RATTUS OR MOUSE OR
MICE OR RODENT#
L4 1 SEA FILE=REGISTRY ABB=ON 9004-34-6
L5 394825 SEA FILE=HCAPLUS ABB=ON L4 OR CORNCOB? OR COB OR COBS OR
MAIZE OR ?CELLULOS?
L31 1658 SEA FILE=WPIX ABB=ON L2 AND (TEST? OR MEASUR? OR SCREEN?) AND
(RODENTIC? OR TOXIC?)

L32 131261 SEA FILE=WPIX ABB=ON L5 OR CORN
 L33 63 SEA FILE=WPIX ABB=ON L31 AND L32
 L34 339776 SEA FILE=WPIX ABB=ON BLOAT? OR IMPACT? OR COMPACT? OR CAECUM
 OR INTESTIN? OR GUT OR STOMACH? OR CECUM
 L35 11 SEA FILE=WPIX ABB=ON L33 AND L34
 L36 2 SEA FILE=WPIX ABB=ON L33 AND (WATER?(2A) (RETEN? OR RETAIN?)
 OR ION(W)TRANSPORT? OR POST MORTEM OR AUTOPS? OR NECROPS?)
 L53 5 SEA FILE=AGRICOLA ABB=ON L35 OR L36
 L54 707 SEA FILE=AGRICOLA ABB=ON RODENTICIDES+NT/CT
 L55 0 SEA FILE=AGRICOLA ABB=ON L54 AND NONTOXIC?
 L56 1 SEA FILE=AGRICOLA ABB=ON L54 AND NON(W)TOXIC?
 L57 66478 SEA FILE=AGRICOLA ABB=ON BLOAT? OR IMPACT? OR COMPACT? OR
 CAECUM OR INTESTIN? OR GUT OR STOMACH? OR CECUM
 L58 21 SEA FILE=AGRICOLA ABB=ON L54 AND L57
 L59 0 SEA FILE=AGRICOLA ABB=ON L58 AND (WATER?(2A) (RETEN? OR
 RETAIN?) OR ION(W)TRANSPORT? OR POST MORTEM OR AUTOPS? OR
 NECROPS?)
 L60 0 SEA FILE=AGRICOLA ABB=ON L58 AND CELLULOS?
 L61 0 SEA FILE=AGRICOLA ABB=ON L58 AND WATER?
 L62 6 SEA FILE=AGRICOLA ABB=ON L53 OR L55 OR L56 OR L59 OR L60 OR
 L61

=> FILE BIOSIS

FILE 'BIOSIS' ENTERED AT 12:08:43 ON 27 MAY 2003
 COPYRIGHT (C) 2003 BIOLOGICAL ABSTRACTS INC. (R)

FILE COVERS 1969 TO DATE.
 CAS REGISTRY NUMBERS AND CHEMICAL NAMES (CNs) PRESENT
 FROM JANUARY 1969 TO DATE.

RECORDS LAST ADDED: 21 May 2003 (20030521/ED)

=> D QUE L70

L2 1457740 SEA FILE=HCAPLUS ABB=ON RAT OR RATS OR RATTUS OR MOUSE OR
 MICE OR RODENT#
 L4 1 SEA FILE=REGISTRY ABB=ON 9004-34-6
 L5 394825 SEA FILE=HCAPLUS ABB=ON L4 OR CORNCOB? OR COB OR COBS OR
 MAIZE OR ?CELLULOS?
 L31 1658 SEA FILE=WPIX ABB=ON L2 AND (TEST? OR MEASUR? OR SCREEN?) AND
 (RODENTIC? OR TOXIC?)
 L32 131261 SEA FILE=WPIX ABB=ON L5 OR CORN
 L33 63 SEA FILE=WPIX ABB=ON L31 AND L32
 L34 339776 SEA FILE=WPIX ABB=ON BLOAT? OR IMPACT? OR COMPACT? OR CAECUM
 OR INTESTIN? OR GUT OR STOMACH? OR CECUM
 L35 11 SEA FILE=WPIX ABB=ON L33 AND L34
 L36 2 SEA FILE=WPIX ABB=ON L33 AND (WATER?(2A) (RETEN? OR RETAIN?)
 OR ION(W)TRANSPORT? OR POST MORTEM OR AUTOPS? OR NECROPS?)
 L63 87 SEA FILE=BIOSIS ABB=ON L35 OR L36
 L64 0 SEA FILE=BIOSIS ABB=ON L63 AND RODENTICID?
 L65 1 SEA FILE=BIOSIS ABB=ON ?CELLULOS? AND RODENTICID?
 L66 0 SEA FILE=BIOSIS ABB=ON RODENT CONTROL+NT/CT
 L67 210 SEA FILE=BIOSIS ABB=ON RODENT(L)CONTROL/IT
 L68 2 SEA FILE=BIOSIS ABB=ON L67 AND ?CELLULOS?
 L69 0 SEA FILE=BIOSIS ABB=ON L63 AND L67
 L70 2 SEA FILE=BIOSIS ABB=ON L64 OR L65 OR L66 OR L68 OR L69

=> DUP REM L30 L38 L62 L70

FILE 'HCAPLUS' ENTERED AT 12:09:04 ON 27 MAY 2003

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2003 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'WPIX' ENTERED AT 12:09:04 ON 27 MAY 2003

COPYRIGHT (C) 2003 THOMSON DERWENT

FILE 'AGRICOLA' ENTERED AT 12:09:04 ON 27 MAY 2003

FILE 'BIOSIS' ENTERED AT 12:09:04 ON 27 MAY 2003

COPYRIGHT (C) 2003 BIOLOGICAL ABSTRACTS INC.(R)

PROCESSING COMPLETED FOR L30

PROCESSING COMPLETED FOR L38

PROCESSING COMPLETED FOR L62

PROCESSING COMPLETED FOR L70

L71 28 DUP REM L30 L38 L62 L70 (2 DUPLICATES REMOVED)

=> D L71 1-28 ALL

L71 ANSWER 1 OF 28 HCAPLUS COPYRIGHT 2003 ACS DUPLICATE 1

AN 2002:555279 HCAPLUS

DN 137:89805

TI **Rodenticide** and **screening** method for
rodenticides

IN Dawson, William

PA Natrocell Technologies Ltd., UK

SO PCT Int. Appl., 30 pp.

CODEN: PIXXD2

DT Patent

LA English

IC ICM A01N065-00

ICS A01N043-16; A01N025-00; G01N037-00; G01N033-50

CC 5-5 (Agrochemical Bioregulators)

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2002056693	A1	20020725	WO 2002-GB171	20020116
	W:				
	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,				
	CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GD, GE, GH, GM,				
	HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,				
	LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL,				
	PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA,				
	UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH,				
	CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR,				
	BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	GB 2374286	A1	20021016	GB 2002-943	20020116
	US 2002160031	A1	20021031	US 2002-46657	20020116
PRAI	GB 2001-1136	A	20010117		

AB A **rodenticide** which is nontoxic to humans, domestic animals and livestock comprises a **water-retentive** material as the active ingredient and a **rodent** attractant. Preferably the **water-retentive** material is **cellulosic** material. In certain embodiments the **water-retentive** material comprises **.alpha.-cellulose**. In one preferred

embodiment the **cellulosic** material comprises purified **cellulose** derived from the core of the **cob** of the DK 446 **maize** hybrid or from the core of the **cob** of an agonist of the DK 446 hybrid. The unique selective **toxicity** of such **rodenticides** arises from their interference with water transport through the **gut** wall, particularly in the **cecum** (where **cellulose** is digested in **rats**). The **cecum** is vestigial in humans, who are not therefore affected by such materials. A lab. method of **screening** candidate **water-retentive** materials for **rodenticidal** activity in the field is disclosed, involving examn. of the **gut** (for **compaction** and **bloating**) and the **cecum** (for **impaction**).

ST **rodenticide corncob cellulose; cecum**
 rodent rodenticide screening

IT **Corn**
 (DK 446 hybrid; **rodenticidal water-retentive cellulose** from **cob** of)

IT **Intestine**
 (cecum, rodent; use for **screening** of **rodenticides**)

IT **Rat** (*Rattus norvegicus*)
 (control by **water-retentive cellulose** from **cob** of DK 446 **maize** hybrid)

IT **Corncob**
 (**rodenticidal water-retentive cellulose** from **cob** of DK 446 **maize** hybrid)

IT **Rodenticides**
 (**water-retentive cellulose** from **cob** of DK 446 **maize** hybrid)

IT **9004-34-6P, Cellulose, biological studies**
 RL: AGR (Agricultural use); BUU (Biological use, unclassified); PUR (Purification or recovery); BIOL (Biological study); PREP (Preparation);
 USES (Uses)

(**rodenticidal water-retentive cellulose** from **cob** of DK 446 **maize** hybrid)

RE.CNT 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD
 RE

- (1) Chuhran, J; WO 9702741 A 1997
- (2) Chuhran, J; WO 9702743 A 1997
- (3) Howard, H; GB 2311464 A 1997 HCAPLUS
- (4) Howard, H; WO 9735470 A 1997

L71 ANSWER 2 OF 28 WPIX (C) 2003 THOMSON DERWENT

AN 2003-140422 [13] WPIX

DNN N2003-111565 DNC C2003-035637

TI Inhibiting tumor growth in mammalian host expressing native stimulatory lectin-like natural killer receptor NKG2D, by administering to host a composition comprising multivalent NKG2D-binding agent.

DC B04 D16 S03

IN DIEFENBACH, A; RAULET, D H

PA (DIEF-I) DIEFENBACH A; (RAUL-I) RAULET D H; (REGC) UNIV CALIFORNIA

CYC 100

PI WO 2002096459 A1 20021205 (200313)* EN 32p A61K039-395

RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW MZ
 NL OA PT SD SE SL SZ TR TZ UG ZM ZW

W: AE AG AL AM AT AU AZ BA BB BG BR BY BZ CA CH CN CO CR CU CZ DE DK
 DM DZ EC EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR

KZ LC LK LR LS LT LU LV MA MD MG MK MN MW MX MZ NO NZ OM PH PL PT
 RO RU SD SE SG SI SK SL TJ TM TN TR TT TZ UA UG UZ VN YU ZA ZM ZW
 US 2002187151 A1 20021212 (200315) A61K039-395
 ADT WO 2002096459 A1 WO 2002-US16924 20020531; US 2002187151 A1 US 2001-871491
 20010531
 PRAI US 2001-871491 20010531
 IC ICM A61K039-395
 ICS A01N061-00; A61K031-00; A61K038-00; C07K001-00; C07K002-00;
 C07K004-00; C07K005-00; C07K007-00; C07K014-00; C07K016-00;
 C07K017-00; G01N033-48
 AB WO 200296459 A UPAB: 20030224

NOVELTY - Inhibiting (M) tumor growth in a mammalian host expressing native stimulatory lectin-like natural killer (NK)G2D receptor and determined to harbor a tumor arising in situ and containing tumor cells or determined to be predisposed to harbor the tumor, comprising administering to the host a composition of a multivalent NKG2D-binding agent, and detecting a resultant inhibition of growth of the tumor, is new.

ACTIVITY - Cytostatic.

The induction of therapeutic response of ligands expressing tumor cells in carcinogen induced tumors was studied. The potential of NKG2D-ligand transduced cells in the treatment of primary cancers induced by chemical carcinogenesis was examined. At the age of 50 days virgin Sprague Dawley rats were given 20 mg dimethylbenzanthracene (DMBA) dissolved at 45 deg. C in 1 ml of corn oil through a stomach catheter. Thereafter, rats were examined to monitor the outgrowth of mammary tumors at weekly intervals for the first 3 weeks, and then twice weekly until the first tumor was detected in each rat. Rats showing signs of persistent toxicity due to DMBA administration (diarrhea, fur ruffling, poor mobility) were excluded from the experiment. Palpable tumors were biopsied and isolated tumor cells were cultured and transduced with NKG2D-ligand expression vectors. At the onset of first palpable tumors, between 40 and 60 days after DMBA administration, rats were allotted to treatment (perfusion of 2 multiply 10⁶ NKG2D-ligand transduced cells) or control. (corresponding untransduced cells) groups, alternating between the two. The treatment period was followed by an observation period of 28 days. Tumors were measured weekly with a caliper and tumor volumes were calculated. A single intragastric dose of 20 mg of DMBA, administered to immature, virgin rats at 50 days of age, was sufficient to induce the onset with 40-50 days of large, fast-growing tumors, localized in the mammary epithelial area, which has been classified as mammary adenocarcinomas. In carcinogen-fed rats, the onset of the first tumor nodule was detectable by palpation as early as 40 days after DMBA administration, followed by other primary tumors appearing within 7-14 days in the mammary area of the rats, reaching the maximum number of five tumors per animal. At the onset of their first palpable tumor, rats were randomly divided into two groups, one group of rats was treated with NKG2D-ligand transduced cells, and a second group of control animals was treated with equivalent untransduced cells. NKG2D-ligand transduced cells showed a powerful inhibitory activity on mammary cancer growth. From the second week of treatment (starting at day 7), up to the end of the off-therapy follow-up period, tumor burden values in the treated group were significantly lower than those in controls. At the end of treatment, the ratio between treated tumor volumes and control tumor volumes (T:C ratios) was less than 0.1. Throughout the whole experiment, no sign of toxicity from the transduced cells was detected.

MECHANISM OF ACTION - Inhibitor of tumor growth (claimed); Enhancer

of immune response to cancer cells.

USE - (M) is useful for inhibiting tumor growth in a mammalian host expressing native NKG2D and determined to harbor a tumor arising in situ and comprising tumor cells or determined to be predisposed to harbor the tumor (claimed). (M) is effective against a wide variety of tumor types such as solid or hematological tumors, melanoma, lymphoma, sarcoma, glioma, leukemia, breast cancer or brain cancer. (M) is useful for treating neoplasia or cancer, or for enhancing the immune response to cancer cells.

Dwg.0/0

FS CPI EPI

FA AB; DCN

MC CPI: B04-G05; B14-H01; B14-H01A; B14-H01B; D05-H09

EPI: S03-E14H

L71 ANSWER 3 OF 28 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.

AN 2002:159699 BIOSIS

DN PREV200200159699

TI Rodent repellent system.

AU Warberg, Kari G. (1)

CS (1) 4551 78th Ave. NW., New Town, ND, 58763-9574 USA

PI US 6337081 January 08, 2002

SO Official Gazette of the United States Patent and Trademark Office Patents, (Jan. 8, 2002) Vol. 1254, No. 2, pp. No Pagination.

<http://www.uspto.gov/web/menu/patdata.html>. e-file.

ISSN: 0098-1133.

DT Patent

LA English

AB A rodent repellent system for repelling rodents within enclosed areas while simultaneously providing a pleasant scent. The inventive device includes a container having an opening, a drawstring within the opening of the container, **cellulose** fiber such as corn cob chips, and a fragrance oil. The fragrance oil preferably has a woody floral blend similar to potpourri. The container is preferably of a perforated material or cloth. A storage box having a lid preferably stores one or more of the containers preferably within a sealable plastic bag. The user attaches the container to a member within the vehicle such as a handle for retaining the container in a prominent position. The fragrance oil is retained by the corn cob chips and slowly released through the container. The fragrance oil provides a strong scent that repels rodents and small animals by irritating their respiratory system while simultaneously providing a pleasant scent to humans.

NCL 424417000

CC Pest Control, General; Pesticides; Herbicides *54600

BC Rodentia - Unspecified 86265

IT Major Concepts

Equipment, Apparatus, Devices and Instrumentation; Pest Assessment
Control and Management

IT Chemicals & Biochemicals

fragrance oil

IT Methods & Equipment

method for controlling rodents: pest **control** method;

rodent repellent system: equipment

ORGN Super Taxa

Rodentia: Mammalia, Vertebrata, Chordata, Animalia

ORGN Organism Name

rodent (Rodentia): pest

ORGN Organism Superterms

Animals; Chordates; Mammals; Nonhuman Mammals; Nonhuman Vertebrates;
Rodents; Vertebrates

L71 ANSWER 4 OF 28 HCAPLUS COPYRIGHT 2003 ACS

AN 2002:348955 HCAPLUS

DN 137:352134

TI Consumption of an omega-3 fatty acids product, INCELL AAFA, reduced
side-effects of CPT-11 (irinotecan) in **mice**

AU Hardman, W. E.; Moyer, M. P.; Cameron, I. L.

CS Pennington Biomedical Research Center, Louisiana State University, Baton
Rouge, LA, 70808, USA

SO British Journal of Cancer (2002), 86(6), 983-988

CODEN: BJCAAI; ISSN: 0007-0920

PB Nature Publishing Group

DT Journal

LA English

CC 18-5 (Animal Nutrition)

AB INCELL AAFA, an omega-3 polyunsatd. fatty acid product contg. a high
concn. of long chain fatty acids, was **tested** for its ability to
ameliorate the harmful side effects of CPT-11 chemotherapy including:
leukopenia, anemia, asthenia, **wt. loss** and liver
involvement. Four groups of **mice** were fed an AIN-76
diet modified to contain: 10% wt./wt. **corn** oil (CO), 0%
AAFA; 9% CO, 1% AAFA; 8% CO, 2% AAFA; or 7% CO, 3% AAFA. After 2 wk on
the **diets**, half of the **mice** received CPT-11
chemotherapy (60 mg kg⁻¹ q 4 days, i.v.) the rest of the **mice**
received vehicle for 2 wk. It was found that 2% AAFA in the **diet**
of the CPT-11 treated **mice**: decreased apoptotic figures in the
duodenal crypts; markedly suppressed the inflammatory eicosanoid,
prostaglandin E2 in the liver; prevented liver hypertrophy; improved white
blood cell counts; significantly increased red blood cell counts;
decreased nos. of CPT-11 induced immature red blood cell and micronuclei
in red blood cells of the peripheral blood; increased eicosapentenoic acid
and docosahexaenoic acid in liver cell membranes and maintained normal
grooming behavior. Thus 2% AAFA in the **diet** reduced the side
effects of CPT-11 treatment in **mice**.

ST omega3 fatty acid INCELL AAFA irinotecan antitumor **toxicity**

IT Antitumor agents

Diet

(effect of omega-3 fatty acids product, INCELL AAFA, on side-effects of
CPT-11 (irinotecan) in **mice**)

IT Fatty acids, biological studies

RL: ADV (Adverse effect, including toxicity); FFD (Food or feed use); PAC
(Pharmacological activity); THU (Therapeutic use); BIOL (Biological
study); USES (Uses)

(polyunsatd., n-3; effect of omega-3 fatty acids product, INCELL AAFA,
on side-effects of CPT-11 (irinotecan) in **mice**)

IT 474453-60-6, AAFA

RL: ADV (Adverse effect, including toxicity); FFD (Food or feed use); PAC
(Pharmacological activity); THU (Therapeutic use); BIOL (Biological
study); USES (Uses)

(effect of omega-3 fatty acids product, INCELL AAFA, on side-effects of
CPT-11 (irinotecan) in **mice**)

IT 100286-90-6, CPT-11

RL: ADV (Adverse effect, including toxicity); THU (Therapeutic use); BIOL
(Biological study); USES (Uses)

(effect of omega-3 fatty acids product, INCELL AAFA, on side-effects of
CPT-11 (irinotecan) in **mice**)

RE.CNT 35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR THIS RECORD
RE

- (1) Anon; Anderson's Pathology 1996, P1793
- (2) Anon; Physicians Desk Reference 2001, P2575
- (3) Anti, M; Gastroenterology 1992, V103, P883 MEDLINE
- (4) Anti, M; Gastroenterology 1994, V107, P1709 MEDLINE
- (5) Barber, M; Br J Cancer 1999, V81, P80 HCAPLUS
- (6) Bartram, H; Gastroenterology 1993, V105, P1317 HCAPLUS
- (7) Burns, C; Clin Cancer Res 1999, V5, P3842
- (8) Calviello, G; Lipids 1999, V34, P599 HCAPLUS
- (9) Das, U; Prostaglandins Leukot Essent Fatty Acids 1998, V58, P39 HCAPLUS
- (10) de Vries, C; Anticancer Res 1992, V12, P1513 HCAPLUS
- (11) Fischer, M; Photochem Photobiol 1991, V54, P381 HCAPLUS
- (12) Germain, E; Lipids 1999, V34, P203
- (13) Hardman, W; Anticancer Res 1999, V19, P2269 HCAPLUS
- (14) Hardman, W; Br J Cancer 1997, V76, P347 HCAPLUS
- (15) Hardman, W; Br J Cancer 1999, V81, P440 HCAPLUS
- (16) Hardman, W; Cancer Lett 2000, V151, P145 HCAPLUS
- (17) Hardman, W; Clin Cancer Res 2001, V7, P2041 HCAPLUS
- (18) Ikuno, N; J Natl Cancer Inst 1995, V87, P1876 HCAPLUS
- (19) Karmali, R; Nutrition 1996, V12, PS2 MEDLINE
- (20) Kase, Y; Jpn J Pharmacol 1997, V75, P399 HCAPLUS
- (21) Meydani, S; J Nutr 1991, V121, P547 HCAPLUS
- (22) Potten, C; Cancer Metastasis Rev 1992, V11, P179 MEDLINE
- (23) Rivory, L; Clin Exp Pharmacol Physiol 1996, V23, P1000 HCAPLUS
- (24) Rothenberg, M; Cancer 1999, V85, P786 HCAPLUS
- (25) Rothenberg, M; J Clin Oncol 1996, V14, P1128 HCAPLUS
- (26) Rothenberg, M; Semin Oncol 1998, V25, P39 HCAPLUS
- (27) Sakai, H; J Physiol 1997, V505.1, P133
- (28) Sakai, H; Naunyn-Schmiedeberg's Arch Pharmacol 1995, V351, P309 HCAPLUS
- (29) Shao, Y; Lipids 1995, V30, P1035 HCAPLUS
- (30) Shao, Y; Nutr Cancer 1997, V28, P63 HCAPLUS
- (31) Slichenmeyer, W; J Natl Cancer Inst 1993, V85, P271
- (32) Takahata, K; Lipids 1999, V34, P105
- (33) Tisdale, M; Prostaglandins Leukot Essent Fatty Acids 1993, V48, P105 HCAPLUS
- (34) Vijayalaxmi; Radiat Res 1997, V147, P495 HCAPLUS
- (35) Workman, P; Br J Cancer 1998, V77, P1

L71 ANSWER 5 OF 28 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved. (2003)

AN 2002:10607 AGRICOLA

DN IND23245672

TI Effect of wheat bran fiber and bran particle size on fat and fiber digestibility and gastrointestinal tract **measurements** in the **rat**.

AU Kahlon, T.S.; Chow, F.I.; Hoefer, J.L.; Betschart, A.A.

SO Cereal chemistry, July/Aug 2001. Vol. 78, No. 4. p. 481-484

Publisher: St. Paul, Minn. : American Association of Cereal Chemists, 1924-

CODEN: CECHAF; ISSN: 0009-0352

NTE Includes references

CY Minnesota; United States

DT Article

FS U.S. Imprints not USDA, Experiment or Extension

LA English

AB The effect of wheat bran (AACC hard red) and bran particle size on fat and fiber digestibility and gastrointestinal tract **measurements** were investigated with diets containing 5.7-10.7% dietary fiber. Fifty-six male weanling Sprague-Dawley **rats** were randomly assigned to four diets containing 5% **cellulose** (C5); 10.5% **cellulose** (C10); 21.5% coarse (2 mm) wheat bran (CB); or 22.2% fine (0.5 mm) wheat bran (FB) in a six-week study. Dietary fiber digestibilities were significantly different ($P < 0.05$) among treatment diets (CB > FB > C5 > C10) but there was no effect in fat digestibility among treatments. High-fiber diets fed to **rats** resulted in significantly greater wet and dry fecal weights than low-fiber diets. Bran diets resulted in significantly higher fecal moisture than **cellulose** diets. **Cecum** lengths increased significantly with bran diets compared with **cellulose** diets. The CB diet resulted in significantly higher **stomach** weights than with **cellulose** diets. **Stomachs** were heavier and cecal lengths were greater with bran diets than with **cellulose** diets; however, a high-**cellulose** diet resulted in increased colon weight. Except for higher fiber digestibility of coarse bran, bran particle size had no significant effects. Healthful effects of wheat bran may be associated with gastrointestinal morphology and function. Fecal bulking and decreased **intestinal** transit time can prevent constipation and may dilute or reduce absorption of **toxic** or carcinogenic metabolites, thus improving gastrointestinal health and lowering the risk of tumor development and cancer.

CC Q504 Food Composition, Field Crop Products; Q104 Food Processing, Field Crop Products; T300 Diet and Diet-related Diseases
CT animal models; **cellulose**; diet; dietary fat; digesta; digestibility; digestive tract; feces; feces composition; fiber; food processing quality; length; moisture content; nutrient intake; particle size; **rats**; weight; wheat bran
RN 9004-34-6 (CELLULOSE)

L71 ANSWER 6 OF 28 HCAPLUS COPYRIGHT 2003 ACS

AN 2001:584478 HCAPLUS

DN 135:288042

TI Subchronic **toxicity** of fish oil concentrates in male and female **rats**

AU Rabbani, P. Isaac; Alam, Hamida Z.; Chirtel, Stuart J.; Duvall, Robert E.; Jackson, Randolph C.; Ruffin, George

CS Risk Assessment, HFS-308, Center for Food Safety and Applied Nutrition, Food and Drug Administration, Washington, DC, 20204, USA

SO Journal of Nutritional Science and Vitaminology (2001), 47(3), 201-212
CODEN: JNSVA5; ISSN: 0301-4800

PB Center for Academic Publications Japan

DT Journal

LA English

CC 18-5 (Animal Nutrition)

Section cross-reference(s): 4

AB The purpose of this study, second in a series, was to evaluate the effects, particularly those that may be harmful, of high-dose, long-term consumption of fish oil concs. (FOC) using male and female **rats**. One hundred and twenty male and 120 female **rats** were gavaged daily with oils and oil mixts. in a vol. equal to 0.5% body wt. (5 mL/kg/d) for 13 wk. The administered oils were **corn** oil, pure menhaden oil (MO), pure MaxEPA fish oil or different mixts. of **corn** oil with MO. The stability and the homogeneity of the dosing solns. were **tested** under study conditions. The animals received

isocaloric and isonitrogenous **diets** throughout. Food and pure water were supplied **ad libitum**. At the end of the in-life phase of the study, the animals were anesthetized with CO₂ and humanely killed by exsanguination. Blood and other tissues were prepd. for various clin., histopathol. and lab. **tests**. Some beneficial effects of FOC, such as redn. in total serum cholesterol, in **rats** were confirmed. However, the authors also obsd. a significant redn. in abs. amt. of serum HDL and a significant increase in relative liver and spleen wts. in both sexes with the high dose of FOC. High doses of FOC (5 mL/kg/d) reduced serum iron and vitamin E concns. A redn. in osmotic fragility of RBC as well as an increase in RBC deformity were also obsd. in **rats** treated with high doses of FOC. These **rats** showed a significant overall increase in WBC count. It can be concluded that in **rats**, subchronic consumption of high levels of FOC can be beneficial but may also be harmful because of induction of clin. abnormalities including increased red cell deformity, increased relative liver and spleen wts., and reduced serum HDL, iron and vitamin E concns.

ST fish oil **diet** subchronic **toxicity**

IT Glycerides, biological studies

RL: BOC (Biological occurrence); BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); OCCU (Occurrence); PROC (Process)

(blood; subchronic **toxicity** of fish oil concs. in male and female **rats**)

IT Fats and Glyceridic oils, biological studies

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(fish, n-3 fatty acid-high, MaxEPA; subchronic **toxicity** of fish oil concs. in male and female **rats**)

IT Fats and Glyceridic oils, biological studies

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(fish; subchronic **toxicity** of fish oil concs. in male and female **rats**)

IT Lipoproteins

RL: BOC (Biological occurrence); BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); OCCU (Occurrence); PROC (Process)

(high-d.; subchronic **toxicity** of fish oil concs. in male and female **rats**)

IT Fats and Glyceridic oils, biological studies

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(menhaden; subchronic **toxicity** of fish oil concs. in male and female **rats**)

IT Fatty acids, biological studies

RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

(polyunsatd., n-3; subchronic **toxicity** of fish oil concs. in male and female **rats**)

IT Body weight

Liver

Reproductive organ

Sex

Spleen

(subchronic **toxicity** of fish oil concs. in male and female **rats**)

IT Mineral elements, biological studies

RL: BOC (Biological occurrence); BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); OCCU (Occurrence); PROC (Process)

(subchronic **toxicity** of fish oil concs. in male and female **rats**)

IT 57-88-5, Cholesterol, biological studies

RL: BOC (Biological occurrence); BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); OCCU (Occurrence); PROC (Process)

(blood; subchronic **toxicity** of fish oil concs. in male and female **rats**)

IT 7439-89-6, Iron, biological studies 7723-14-0, Phosphorus, biological studies 9000-86-6, Glutamic pyruvic transaminase 9000-97-9 9001-60-9, Lactic dehydrogenase 9001-78-9, Alkaline phosphatase 16887-00-6, Chloride, biological studies 189764-17-8, Vitamin E

RL: BOC (Biological occurrence); BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); OCCU (Occurrence); PROC (Process)

(subchronic **toxicity** of fish oil concs. in male and female **rats**)

RE.CNT 69 THERE ARE 69 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Abramowicz, M; Medical Lett 1998, V40, P117
- (2) Anderson, R; Cancer Lett 1987, V36, P55 MEDLINE
- (3) Anon; Official Methods of Analysis of the Association of Official Analytical Chemists, 15th ed 1990, VII(sec 965.33), P956
- (4) Bartoli, G; Bioch Bioph Res Com 1995, V213, P881 HCAPLUS
- (5) Bellisola, G; Clin Chem Acta 1992, V205, P75 HCAPLUS
- (6) Berdanier, C; J Nutr 1992, V122, P1309 HCAPLUS
- (7) Berlin, E; Comp Biochem Physiol 1994, V109A, P53 HCAPLUS
- (8) Borge, N; Cardiology 1998, V89, P94
- (9) Byleveld, P; J Nutr 1999, V129, P328 HCAPLUS
- (10) Chang, H; Metabolism 1992, V41, P1 HCAPLUS
- (11) Cho, S; J Nutr Sci Vitaminol 1995, V41, P61 HCAPLUS
- (12) Choi, Y; South Med J 1998, V91, P372 MEDLINE
- (13) Dacie, J; Protocol Hematology, 4th ed 1968, P166
- (14) De Craemer, D; Eur J Morphol 1993, V31, P77 MEDLINE
- (15) Einig, R; J Am Oil Chem 1987, V64, P499
- (16) Ershoff, B; J Nutr 1980, V71, P45
- (17) Everett, R; The Mouse in Biomedical Research 1983, P313
- (18) Falton, C; Lancet 1994, V344, P1195
- (19) Farwer, S; Br J Nutr 1994, V72, P127 HCAPLUS
- (20) Foulon, T; Scand J Clin Lab Invest 1999, V59, P239 HCAPLUS
- (21) Fowler, J; Arch Toxicol Suppl 1982, V5, P152 MEDLINE
- (22) Graham, R; J Histochem Cytochem 1966, V14, P291 HCAPLUS
- (23) Halminski, M; J Nutr 1991, V121, P1554 HCAPLUS
- (24) Harris, W; Clin Cardiol 1999, V22(6 Suppl), P1140 MEDLINE
- (25) Hartog, J; Am J Clin Nutr 1987, V46, P258 HCAPLUS
- (26) Herold, M; Am J Clin Nutr 1986, V43, P566
- (27) Hubbard, N; Cancer Lett 1998, V124, P1 HCAPLUS
- (28) Huff, M; Biochim Biophys Acta 1993, V1210, P113 HCAPLUS
- (29) Kinsella, J; Nutr Today 1986, V21, P7
- (30) Kirchgessner, M; Z Ernahrungswiss 1994, V33, P146 HCAPLUS
- (31) Kirmani, Z; J Burn Care Rehabil 1995, V16, P306 MEDLINE
- (32) Knapp, H; J Am Coll Nutr 1990, V9, P344 MEDLINE
- (33) Kolanowski, W; Int J Food Sci Nutr 1999, V50, P39 HCAPLUS
- (34) Kremer, J; Am J Clin Nutr 2000, V71, P349S HCAPLUS
- (35) Kromann, N; Acta Med Scand 1980, V208, P401 MEDLINE

- (36) Kuralay, F; J Toxcol Environ Health A 1998, V53, P223 HCAPLUS
- (37) Leblanc, M; Am J Kidney Dis 1998, V32(3 Suppl), P444
- (38) Loeb, W; The Clinical Chemistry of Laboratory Animals 1989
- (39) Malasanos, T; Diabetes Care 1991, V14, P1160 MEDLINE
- (40) Mitruka, B; Clinical Biochemical and Hematological Reference Values in Normal Experimental Animals 1977
- (41) Mori, T; Am J Clin Nutr 1994, V59, P1060 HCAPLUS
- (42) Morimoto, M; Proc Soc Exp Biol Med 1994, V207, P197 HCAPLUS
- (43) Muggli, R; Health Effects of Fish and Fish Oil 1989, P201
- (44) Nelson, G; Lipids 1997, V32, P1129 HCAPLUS
- (45) Nieuwenhuys, C; Arterioscler Thromb Vasc Biol 1998, V18, P1480 HCAPLUS
- (46) Oarada, M; Biochim Biophys Acta 2000, V1487, P1 HCAPLUS
- (47) Parks, J; J Lipid Res 1991, V32, P305 HCAPLUS
- (48) Peck, M; New Horiz 1994, V2, P230 MEDLINE
- (49) Perez-Granados, A; Analyst 1997, V120, P899
- (50) Pownall, H; Atherosclerosis 1999, V143, P285 HCAPLUS
- (51) Prasad, A; Am J Med 1961, V31, P532 MEDLINE
- (52) Rabbani, P; J Nutr Sci Vitaminol 1999, V45, P553 HCAPLUS
- (53) Rogers, K; Am J Path 1988, V132, P382 MEDLINE
- (54) Rogers, K; Am J Pathol 1990, V137, P945 HCAPLUS
- (55) SAS Institute; SAS User's Guide: Statistics, 5th Ed 1985
- (56) Semplicini, A; Pharmacol Ther 1994, V61, P385 HCAPLUS
- (57) Smit, M; J Lipid Res 1994, V35, P301 HCAPLUS
- (58) Stone, N; Circulation 1996, V94, P2337 MEDLINE
- (59) Surette, M; J Lipid Res 1992, V33, P263
- (60) Tatenos, S; Nephron 1997, V77, P86 HCAPLUS
- (61) Thomas, T; J Appl Physiol 2000, V88, P2199 HCAPLUS
- (62) Tripodi, A; Biochim Biophys Acta 1991, V1083, P298 HCAPLUS
- (63) Tsai, C; Nutr Res 1989, V9, P673
- (64) Van Elswyk, M; Poult Sci 1994, V73, P653 HCAPLUS
- (65) Villegas, A; Hematologica 1998, V83, P99 MEDLINE
- (66) Vognild, E; Lipids 1998, V33, P427 HCAPLUS
- (67) Wander, R; J Nutr 1997, V127, P1198 HCAPLUS
- (68) Wickwire, K; Proc Soc Exp Biol Med 1995, V9, P397
- (69) Ziboh, V; Am J Clin Nutr 2000, V71, P361S HCAPLUS

L71 ANSWER 7 OF 28 HCAPLUS COPYRIGHT 2003 ACS

AN 1999:5293 HCAPLUS

DN 130:193089

TI Susceptibility of roof-rats (*Rattus rattus*)
to anticoagulant **rodenticides** baits in comparison with
Rattus norvegicus var. *albus*

AU Youssef, Hamada M.; El-Sebai, Mohamed A.; Desheesh, Mohamed A.

CS Department of Agricultural Animal Pests, Plant Protection Research
Institute, Alexandria, Egypt

SO Alexandria Science Exchange (1998), 19(3), 497-506
CODEN: ALSEEF; ISSN: 1110-0176

PB Prof. Dr. A. M. Balba Group for Soil and Water Research

DT Journal

LA English

CC 5-5 (Agrochemical Bioregulators)

AB Susceptibility of wild roof rats (*Rattus
rattus*) and albino Norway rats to anticoagulant
rodenticides with different formulated baits were evaluated using
lab. **feeding free choice test**. Biochem.
measurements (Hb, haematocrit, RBC's, WBC's values as well as
sGOT, sGPT activities in addn. to total serum protein values) in both
strains of **rats** were detd. The wild roof **rats** were

found to be less susceptible to coumatetralyl (0.0375% crushed maize and 0.045% wheat baits) than the albino Norway rats, due to the accumulation of higher dosages of active ingredient more than white rats. Chlorophacinone (0.005% crushed maize bait) was more active against both strains of rats. Hb, haematocrit and RBC's were slightly affected with the tested anticoagulant. While WBC's and sGPT were highly elevated in the blood of *Rattus rattus*. On the other hand, sGOT activities were moderately reduced in the two strains except in case of chlorophacinone which weakly elevated sGOT in *Rattus rattus*. Total serum protein was significantly reduced by the three tested anticoagulants in both strains of rats.

- ST anticoagulant **rodenticide rat** susceptibility;
Rattus susceptibility anticoagulant **rodenticide**
- IT Erythrocyte
 Hematocrit
 Leukocyte
 (anticoagulant **rodenticides** baits effect on *Rattus rattus* and *Rattus norvegicus* var. albus)
- IT Hemoglobins
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
 (anticoagulant **rodenticides** baits effect on *Rattus rattus* and *Rattus norvegicus* var. albus)
- IT **Rodenticides**
 (blood coagulation-inhibiting; susceptibility of roof-rats to anticoagulant **rodenticides** baits in comparison with *Rattus norvegicus* var. albus)
- IT Proteins, general, biological studies
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
 (blood; anticoagulant **rodenticides** baits effect on *Rattus rattus* and *Rattus norvegicus* var. albus)
- IT Rat (*Rattus norvegicus albinus*)
 Rat (*Rattus rattus*)
 (susceptibility of roof-rats to anticoagulant **rodenticides** baits in comparison with *Rattus norvegicus* var. albus)
- IT 9000-86-6, GPT 9000-97-9, GOT
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
 (anticoagulant **rodenticides** baits effect on *Rattus rattus* and *Rattus norvegicus* var. albus)
- IT 3691-35-8, Kaide 5836-29-3, Racumin
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)
 (susceptibility of roof-rats to anticoagulant **rodenticides** baits in comparison with *Rattus norvegicus* var. albus)

RE.CNT 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS RECORD

RE

- (1) Ahmed, N; Ass J of Agricul Sci 1990, V21(4), P19
- (2) Bell, G; Text book of physiology and biochemistry, 9th ed 1976, P246
- (3) Brooks, J; Laboratory evaluation of rodenticides for use against South-Cost Asian Commensal small mammals 1979, V179-720(1-15)
- (4) Dacie, J; Practical haematology 7th ed 1991, P629
- (5) Desheesh, M; J Pest Control & Environ Sci 1991, V3(1), P102

- (6) Desheesh, M; Proc 4th Pesticides Conf 1981, VIII
- (7) Hadler, M; J Hyg Camb 1975, V74, P441 HCAPLUS
- (8) Kumar, D; C F B A 1992, V94(2), P23457
- (9) Kumar, D; Indian J Comp Anim Physiol 1991, V9(2), P81
- (10) Mukatha, K; Pestic Sci 1978, V9, P44
- (11) Reitman, S; Amer J Clin Path 1957, V28, P56 HCAPLUS
- (12) Rowe, F; J Hyg Camb 1965, V63, P417 HCAPLUS
- (13) Steel, R; "Principles and procedures of statistics A Biometrical Approach"
2nd 1980, P633
- (14) Wintrobe, M; Clinical haematology 4th ed 1965

L71 ANSWER 8 OF 28 AGRICOLA Compiled and distributed by the National
Agricultural Library of the Department of Agriculture of the United States
of America. It contains copyrighted materials. All rights reserved.

(2003)

AN 1998:31121 AGRICOLA

DN IND20815103

TI Bait placement and acceptance by rats in macadamia orchards.

AU Tobin, M.E.; Sugihara, R.T.; Koehler, A.E.

AV DNAL (SB599.C8)

SO Crop protection, Sept 1997. Vol. 16, No. 6. p. 507-510

Publisher: Kidlington, Oxford, UK : Elsevier Science Ltd.

CODEN: CRPTD6; ISSN: 0261-2194

NTE Includes references

CY England; United Kingdom

DT Article

FS Non-U.S. Imprint other than FAO

LA English

AB Black rats (*Rattus rattus*) cause extensive damage in Hawaiian macadamia
(*Macadamia integrifolia*) orchards. Many growers apply rodenticides to
reduce rat populations in orchards, but improper placement of bait may
reduce the effectiveness of many baiting programs. We evaluated the
optimum placement of bait in macadamia orchards among the three locations
specified on current rodenticide labels. We placed a **non-**
toxic oat bait treated with 0.75% tetracycline hydrochloride, an
antibiotic that chelates with calcium in growing bones and teeth and
fluoresces under UV light, in burrows, on the ground and in trees in
separate orchard sections. We consistently captured the greatest
percentage of marked rats (53-91%) in sections where we placed the bait in
trees and the lowest percentage of marked rats (0-36%) where we broadcast
bait on the ground. Placement of bait in burrows produced intermediate
results (40-70%). These results suggest that broadcasting bait on the
orchard floor reduces the effectiveness of rat control programs. Placing
baits in trees targets rats that not only are most likely to eat the
poison bait, but also are most likely to damage developing nuts.

CC F820 Pests of Plants, Animals; L300 Animal Ecology and Behavior

CT activity; baiting; baits; macadamia integrifolia; movement; placement;
rattus rattus; rodent control; **rodenticides**; vertebrate pests

ST animal activity; optimum location

GT hawaii

RN 64-75-5 (TETRACYCLINE HYDROCHLORIDE)

7440-70-2 (CALCIUM)

L71 ANSWER 9 OF 28 HCAPLUS COPYRIGHT 2003 ACS

AN 1997:627605 HCAPLUS

DN 127:244083

TI Acute pathological and biochemical effects of 2,4,6-trinitrophenyl-N-
methylnitramine (tetryl) in male **rats**

AU Reddy, Gunda; Qualls, C. W., Jr.; Hampton, A. E. G.; Yelamanchili, A.; Kim, S.
CS U.S. Army Center for Health Promotion and Preventive Medicine, Aberdeen Proving Ground, MD, 21010-5422, USA
SO Research Communications in Pharmacology and Toxicology (1997), 2(1&2), 1-11
CODEN: RCPTFY; ISSN: 1087-1101
PB PJD Publications
DT Journal
LA English
CC 4-3 (Toxicology)
AB Male Fischer 344 **rats** were gavaged with 2,4,6-trinitrophenyl-N-methylnitramine (tetryl) in **corn** oil at 0, 500, or 1000 mg/kg. Twenty-four hours after a single treatment, hepatic and **testicular** drug-metabolizing enzymes, hematol., and histopathol. changes were studied. Tetryl did not produce changes in hepatic microsomal cytochrome P 450 and cytochrome b5 content but it significantly increased ethoxy and pentoxy O-dealkylase activities. Cytosolic glutathione S-transferase activity and the content of oxidized or reduced glutathione were not effected in the liver and **testis**. Hematol. parameters altered included an increase in the metHb, glucose, and serum urea nitrogen content and a decrease in lymphocytes. Gross **necropsy** revealed the presence of food contg. yellowish **test** substance and dark brown to red blood deposits in the **stomach** of treated **rats** even though they had been fasted for 16 h. Histopathol. changes obsd. were glycogen accumulation in the liver and focal coagulative necrosis of the gastric mucosa at the junction of the **stomach** and duodenum of treated **rats**. No other treatment-related changes were obsd. in gross or histol. evaluation.
ST tetryl acute **toxicity** biochem
IT Liver
Lymphocyte
Microsome
Stomach
Testis
(acute pathol. and biochem. effects of tetryl)
IT Hemoglobins, methemoglobins
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
(acute pathol. and biochem. effects of tetryl)
IT Cytoplasm
(cytosol; acute pathol. and biochem. effects of tetryl)
IT Enzymes, biological studies
RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
(drug-metabolizing; acute pathol. and biochem. effects of tetryl)
IT **Stomach**
(mucosa; acute pathol. and biochem. effects of tetryl)
IT 479-45-8, Tetryl
RL: ADV (Adverse effect, including toxicity); BIOL (Biological study)
(acute pathol. and biochem. effects of tetryl)
IT 9035-39-6, Cytochrome b5 9035-51-2, Cytochrome P 450, biological studies
50812-37-8, Glutathione S-transferase 59793-97-4, Ethoxyresorufin O-dealkylase 96595-04-9, Pentoxyresorufin O-dealkylase
RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)

- (acute pathol. and biochem. effects of tetryl)
- IT 50-99-7, Glucose, biological studies 57-13-6, Urea, biological studies 70-18-8, GSH, biological studies 27025-41-8, GSSG
RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
(acute pathol. and biochem. effects of tetryl)
- L71 ANSWER 10 OF 28 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved.
(2003)
- AN 94:73305 AGRICOLA
DN IND20421203
TI Subchronic **toxicity** studies of SALATRIM structured triacylglycerols in **rats**. 2. Triacylglycerols composed of stearate, acetate, and propionate.
AU Hayes, J.R.; Wilson, N.H.; Pence, D.H.; Williams, K.D.
AV DNAL (381 J8223)
SO Journal of agricultural and food chemistry, Feb 1994. Vol. 42, No. 2. p. 539-551
Publisher: Washington, D.C. : American Chemical Society.
CODEN: JAFCAU; ISSN: 0021-8561
- NTE Includes references
CY District of Columbia; United States
DT Article
FS U.S. Imprints not USDA, Experiment or Extension
LA English
AB SALATRIM 23CA lot A014, SALATRIM 32CA lot A015, and SALATRIM 23SO lot A026 are members of a family of structured triacylglycerols having caloric densities of 4.5-6.0 kcal/g. **Rats** received 0%, 2%, 5% and 10% of the first two SALATRIM fats or 10% **corn** oil in the diet for 13 weeks. Body weight and feed consumption were unaffected. Minimally increased urinary phosphorus clearance occurred in 10% SALATRIM groups. Bone mineral variations and an increased incidence of renal mineralization occurred in 10% SALATRIM and **corn** oil fed **rats**. These changes appeared to be directly related to the quantity of unsaturated fatty acids in the high-fat diets. **Measurements** of serum and liver fat-soluble vitamin concentrations, **necropsy**, clinical pathology, organ weights, and histopathology revealed no SALATRIM-related effects. **Corn** oil produced hepatocellular vacuolation. In a supplementary short-term study, 10% dietary SALATRIM 23SO lot A026 produced no effect on selected serum transaminases. Overall, the SALATRIM fats produced no **toxicologically** significant effects.
- CC Q500 Food Composition and Quality; T300 Diet and Diet-related Diseases
CT acetic acid; dietary fat; lipid substitutes; low fat products; **maize** oil; propionic acid; stearic acid; **toxicity**; triacylglycerols
- RN 57-11-4 (STEARIC ACID)
64-19-7 (ACETIC ACID)
79-09-4 (PROPIONIC ACID)
7723-14-0 (PHOSPHORUS)
8001-30-7 (CORN OIL)
64706-27-0 (SALATRIM 23CA)
64706-27-0 (SALATRIM 23SO)
64706-27-0 (SALATRIM 32CA)
- L71 ANSWER 11 OF 28 HCAPLUS COPYRIGHT 2003 ACS
AN 1994:402906 HCAPLUS

DN 121:2906
TI Ten and ninety-day **toxicity** studies of chloropicrin in
Sprague-Dawley **Rats**
AU Condie, L. W.; Daniel, F. Bernard; Olson, Greg R.; Robinson, Merrel
CS U.S. Army Dugway Proving Ground, Dugway, UT, 84022, USA
SO Drug and Chemical Toxicology (1977) (1994), 17(2), 125-37
CODEN: DCTODJ; ISSN: 0148-0545
DT Journal
LA English
CC 4-3 (Toxicology)
AB The **toxicity** of chloropicrin (CP) was assessed following its
administration to **rats** via oral gavage for either 10 or 90
consecutive days at dose levels of 10, 20, 40, and 80 mg/kg and 2.8 and 32
mg/kg, resp. Control **rats** received **corn** oil at a dose
of 1.0 mL/kg. **Toxicol.** observations include organ and body wt.
measurements, necropsy and histopathol. observations,
urinalysis, clin. chem. and hematol. detns. The most remarkable
toxicol. finding in both studies was the corrosive property of CP
on forestomach tissue. Inflammation, necrosis, acantholysis,
hyperkeratosis and epithelial hyperplasia of the forestomach were seen in
all dose groups of the 10-day study. Similar changes were detected in
only the high dose group in the 90-day study. Decreased red blood cell
parameters were noted in the highest dose groups in both studies, possibly
due to blood loss via the damaged **stomach** lining. CP may have
been aspirated into the lungs of animals in the high dose group in the
ninety day study resulting in pulmonary complications leading to the
deaths of 60% of the males and 80% of the the females starting at week
five. The 8 mg/kg dose group in the ninety day study was considered to be
the no obsd. adverse effect level.
ST **toxicity** chloropicrin
IT Erythrocyte
Hematocrit
Leukocyte
Reticulocyte
Hemoglobins
RL: BIOL (Biological study)
(chloropicrin effect on)
IT Animal growth
(chloropicrin effect on, **toxicity** in relation to)
IT Sex
(chloropicrin **toxicity** in relation to)
IT Organ
(wt. of, chloropicrin effect on)
IT **Stomach, toxic** chemical and physical damage
(forestomach, chloropicrin **toxicity** to)
IT 76-06-2, Chloropicrin
RL: ADV (Adverse effect, including toxicity); BIOL (Biological study)
(**toxicity** of)
L71 ANSWER 12 OF 28 AGRICOLA Compiled and distributed by the National
Agricultural Library of the Department of Agriculture of the United States
of America. It contains copyrighted materials. All rights reserved.
(2003)
AN 1998:19151 AGRICOLA
DN IND20621181
TI Developmental effects of an environmental antiandrogen: the fungicide
vinclozolin alters sex differentiation on the male **rat**.
AU Gray, L.E. Jr; Ostby, J.S.; Kelce, W.R.

CS United State Environmental Protection Agency, Research Triangle Park, NC.
SO Toxicology and applied pharmacology, Nov 1994. Vol. 129, No. 1. p. 46-52
Publisher: Orlando, Fla. : Academic Press Inc.
CODEN: TXAPA9; ISSN: 0041-008X

NTE Includes references

CY Florida; United States

DT Article

FS U.S. Imprints not USDA, Experiment or Extension

LA English

AB In humans and **rodents**, exposure to hormonally active chemicals during sex differentiation can produce a wide range of abnormal sexual phenotypes including masculinized and defeminized females and feminized and demasculinized males. Although numerous "environmental estrogens," including pesticides, **toxic** substances (PCBs), and plant and fungal estrogens, have been shown to alter mammalian sex differentiation, similar information on environmental androgens is lacking. Recently, the fungicide vinclozolin (V) was found to inhibit sexual differentiation in male **rats** in an antiandrogenic manner. In the present study, V was administered to pregnant **rats** (po) at 0, 100, or 200 mg/kg/day in **corn** oil during the period of sex differentiation (Gestational Day 14 to Postnatal Day 3) to examine the demasculinizing effect of this fungicide more closely. In both groups of V-treated male offspring, anogenital distance was female like at birth, and nipple development was prominent at 2 weeks of age. After puberty, most of the V-treated male offspring were unable to attain intromission even though they all mounted sexually receptive females. The V-treated male offspring that appeared to achieve intromission, failed to ejaculate normally, as no sperm were found in the uterus after overnight matings. A factor in the abnormal ejaculation was that all V-treated male offspring had cleft phallus with hypospadias. In addition, a number of unusual reproductive malformations were noted when the males were **necropsied** at 1 year. Many V-treated male offspring had suprainguinal ectopic scrota/**testes**, a vaginal pouch, epididymal granulomas, and small to absent sex accessory glands. During the study, about 25% of the V-treated males died as a result of bladder stones, hydroureter, or hydronephrosis, while other males displayed these lesions at **necropsy**. While some of the above malformations in male offspring can also be produced by perinatal administration of a potent estrogen, like DES, V-treated female offspring did not display any estrogen-like alterations of reproductive development or fecundity. The only change seen in the female offspring was a reduced anogenital distance during neonatal life. Our observation of perinatal-induced agenesis of the prostate and blocked **testicular** descent, a pattern of malformations nearly identical to that reported for the antiandrogen flutamide, is consistent with other recent evidence that this fungicide is an androgen-receptor antagonist.

CC H000 Pesticides, General; X380 Human Medicine, Health and Safety
CT accessory glands; ejaculation; epididymis; intromission; male animals; malformations; penis; pregnancy; progeny; **rats**; scrotum; sex differentiation; sex differentiation disorders; teratogenesis; **testes**; **toxicity**; vinclozolin

ST phallus

RN 8001-30-7 (CORN OIL)

13311-84-7 (FLUTAMIDE)

50471-44-8 (VINCLOZOLIN)

L71 ANSWER 13 OF 28 HCAPLUS COPYRIGHT 2003 ACS

AN 1991:466549 HCAPLUS

DN 115:66549

- TI Biological and chemical characterization of metabolites of *Fusarium moniliforme* isolates
AU Hassanin, Nagawa; Gabal, M. A.
CS Dep. Nutr. Food Chem., Ain-Shams Univ., Egypt
SO Veterinary and Human Toxicology (1990), 32(6), 536-40
CODEN: VHTODE; ISSN: 0145-6296
DT Journal
LA English
CC 4-5 (Toxicology)
Section cross-reference(s): 10, 17
- AB Metabolites of *F. moniliforme* isolates from different types of **corn** were characterized biol. and chem. The biol. assays included **rat feeding, rat dermal tests**, and inoculation of embryonated eggs. The metabolites were identified as diacetoxyscirpenol, ipomeanol, ipomeanine, and diplodiatoxin. The biol. **tests** revealed significant **wt. loss** in **rats** fed the contaminated **corn** for 5 wk. Hemorrhages and edema in the brain and **intestine** were detected in all the **rats**.
- ST *Fusarium* metabolite **toxicity**
IT **Toxicity**
(of *Fusarium moniliforme* metabolites)
IT Mycotoxins
RL: BIOL (Biological study)
(*Fusarium moniliforme* metabolites)
IT 496-06-0, Ipomeanine 2270-40-8 41060-01-9, Diplodiatoxin
135014-96-9, 3-Ipomeanol
RL: PROC (Process)
(as metabolite of *Fusarium moniliforme*, biol. and chem. characterization of)
- L71 ANSWER 14 OF 28 HCAPLUS COPYRIGHT 2003 ACS DUPLICATE 2
AN 1990:135702 HCAPLUS
DN 112:135702
- TI Absence of trichothecenes in toxigenic isolates of *Fusarium moniliforme*
AU Mirocha, C. J.; Abbas, H. K.; Vesonder, R. F.
CS Dep. Plant Pathol., Univ. Minnesota, St. Paul, MN, 55108, USA
SO Applied and Environmental Microbiology (1990), 56(2), 520-5
CODEN: AEMIDF; ISSN: 0099-2240
DT Journal
LA English
CC 10-1 (Microbial Biochemistry)
- AB Thirty-four isolates of *F. moniliforme* were obtained from cereal grains collected in various parts of the world. The isolates were grown on rice and **tested** as a **diet** for **toxicity** to **rats**. Of these isolates, 53% caused death, 12% caused congestion and hemorrhage of the **stomach** and **intestine** as well as hematuria, 21% caused diarrhea, 38% caused **wt. loss**, and 9% were nontoxic. The cultures were **tested** for T-2, HT-2, neosolaniol, acetyl-T-2, T-2-tetraol, iso-T-2, diacetoxyscirpenol, monoacetoxyscirpenol, deoxynivalenol, nivalenol, fusarenone-X, 3-acetyldeoxynivalenol, 15-acetyldeoxynivalenol, zearalenone, moniliformin, fusarochromanone, fusarin-C, and wortmannin; all were neg. In addn., *F. moniliforme* NRRL A25820 was grown on **corn** and banana fruit as solid substrates as well as on a defined liq. medium; none of the above toxins were found. When *F. moniliforme* NRRL A25820 was incorporated into a **rat diet**, **no toxicity** was noted. Twenty-eight addnl. isolates of *E. moniliforme*, isolated from

feed assocd. with equine leukoencephalomalacia, were grown on cracked **corn** for 2 wk. The cultures were neg. when tested for deoxynivalenol, 15-acetyldeoxynivalenol, diacetoxyscirpenol, monoacetoxyscirpenol, nivalenol, and fusarenone X. Seventy-five percent of the isolates were **toxic** to ducklings, indicating the presence of a toxin other than trichothecenes. The results support the conclusion that *F. moniliforme* does not produce trichothecenes.

ST *Fusarium toxigenic* trichothecene absence

IT *Fusarium moniliforme*

(toxigenic, absence of trichothecenes in)

IT Sesquiterpenes and Sesquiterpenoids

RL: BIOL (Biological study)

(trichothecane, absence of, in toxigenic *Fusarium moniliforme*)

L71 ANSWER 15 OF 28 AGRICOLA Compiled and distributed by the National Agricultural Library of the Department of Agriculture of the United States of America. It contains copyrighted materials. All rights reserved. (2003)

AN 89:69538 AGRICOLA

DN IND89035967

TI Utilization of four cultivars of grain amaranth growth in **rats**.

AU Pond, W.G.; Lehmann, J.; Clark, R.

CS USDA, ARS, Roman L. Hruska U.S. Meat Animal Research Center, Clay Center, NE

AV DNAL (RC620.A1N8)

SO Nutrition reports international, May 1989. Vol. 39, No. 5. p. 1081-1089

Publisher: Stoneham, Mass. : Butterworth.

CODEN: NURIBL; ISSN: 0029-6635

NTE Includes references.

DT Article

FS U.S. Imprints not USDA, Experiment or Extension

LA English

AB Growing Sprague-Dawley male **rats** were used in two experiments to determine the nutritional adequacy of four cultivars of grain amaranth for growth. The amaranth cultivars (16.8, 16.0, 16.4 and 16.1% protein) were fed as the sole source of protein and energy or diluted with **maize** to provide 46.1% amaranth in the diet. All cultivars were fed in the ground, unheated form. A diet containing **maize** as the sole source of protein (10.3%) and energy was used as a control and a 16% protein **maize**-soybean meal diet was fed as a reference diet. **Rats** were fed ad libitum for 14 days in individual wire-bottom cages in a light- and temperature-controlled room. In each experiment (5 **rats** fed each of 10 diets in Exp. 1 and 10 **rats** fed each of 6 diets in Exp. 2), body weight gain, feed intake and gain to feed ratio were recorded for each **rat**. Three cultivars (*Amaranthus hypochondriacus* 1024, *A. hypochondriacus* 1046 and *A. hypochondriacus* K188) produced weight gain and feed utilization significantly greater ($P < 0.01$) than **maize**; two cultivars (*A. hypochondriacus* 1046 and *A. hypochondriacus* 1046 and K 188) produced weight gain not significantly different from that obtained with the **maize**-soy reference diet, although daily feed intake was greater with the reference diet. One cultivar (*A. cruentus* 1011) allowed normal growth during week 1, followed by steady weight loss and decreased feed intake during week 2. Refeeding for 7 days of half the **rats** fed 1011 with **maize**-soy reference diet resulted in rapid weight recovery and absence of gross pathology of liver, kidney, **stomach**, spleen, adrenal, and **testes** at slaughter. The nature of the **toxic** factor

present in *A. cruentus* 1011 is unknown, but ingestion seems not associated with permanent organ damage after 2 weeks of feeding. We conclude that three of the four amaranth cultivars **tested** promote growth of **rats** superior to that obtained with **maize** and comparable to that obtained with a 16% **maize**-soy diet.

CC Q504 Food Composition, Field Crop Products; F200 Plant Breeding and Genetics

CT *amaranthus caudatus*; *amaranthus leucocarpus*; cultivars; nutritional value; **rats**

RN 68308-36-1 (SOYBEAN MEAL)

L71 ANSWER 16 OF 28. HCAPLUS COPYRIGHT 2003 ACS

AN 1989:422330 HCAPLUS

DN 111:22330

TI Hepatotoxicity and renal **toxicity** in **rats** of **corn** samples associated with field cases of equine leukoencephalomalacia

AU Voss, K. A.; Norred, W. P.; Plattner, R. D.; Bacon, C. W.

CS Richard B. Russell Agric. Res. Cent., Agric. Res. Serv., Athens, GA, 30613, USA

SO Food and Chemical Toxicology (1989), 27(2), 89-96

CODEN: FCTOD7; ISSN: 0278-6915

DT Journal

LA English

CC 17-5 (Food and Feed Chemistry)

AB A bioassay was developed to det. the potential **toxicity** of **corn** naturally contaminated with *Fusarium moniliforme*. Two groups of five male Sprague-Dawley **rats** were each fed one of two *F. moniliforme*-contaminated **corn** samples, designated CS-1 and CS-2, that were assocd. with sep. field cases of equine leukoencephalomalacia. A control group, also consisting of 5 male **rats**, was fed uncontaminated seed **corn**. All animals survived to the end of the study, and there were no apparent differences in appearance or behavior among groups. **Wt. loss** and irregular food consumption occurred in all groups and probably resulted from nutritional deficiencies inherent in the **corn diets**. Hepatocellular degeneration, necrosis and hyperplasia as well as biliary hyperplasia were found in the **test** groups only and were attributed to *F. moniliforme*. Serum transaminase and alk. phosphatase activities in animals fed CS-1 and CS-2 for 4 wk were significantly increased compared with the controls, while serum bilirubin concn. was increased only in the CS-1 group. Tubular nephrosis was also present in the renal cortex of all animals fed CS-1 and CS-2. These effects may have been related to fumonisins B1 and B2, recently discovered metabolites of *F. moniliforme*, that were found in both CS-1 and CS-2. Short-term studies of this type may be useful in **screening** naturally-contaminated grains and other materials for hepatotoxic metabolites produced by *F. moniliforme*.

ST **corn** hepatotoxicity *Fusarium* toxin leukoencephalomalacia; kidney **corn** *Fusarium* toxin leukoencephalomalacia; fumonisin **corn** bioassay

IT **Feed** contamination

(by hepatotoxins, of **corn**, bioassay of)

IT Kidney, **toxic** chemical and physical damage

Liver, **toxic** chemical and physical damage

(**corn** contaminated with *Fusarium moniliforme* **toxicity** to, equine leukoencephalomalacia in relation to)

IT *Fusarium moniliforme*


- (hepatotoxicity and renal **toxicity** of **corn** contaminated with, equine leukoencephalomalacia in relation to)
- IT **Corn**
(hepatotoxicity and renal **toxicity** of Fusarium-contaminated, equine leukoencephalomalacia in relation to)
- IT **Horse**
(leukoencephalomalacia of, **dietary** Fusarium-contaminated **corn** assocd. with, hepatotoxicity and renal **toxicity** in relation to)
- IT **Brain, disease or disorder**
(leukoencephalomalacia, equine, **dietary** Fusarium-contaminated **corn** assocd. with, hepatotoxicity and renal **toxicity** in relation to)
- IT 635-65-4, Bilirubin, biological studies 9001-78-9, Alkaline phosphatase
RL: BIOL (Biological study)
(of blood serum, increase of, by **dietary corn** contaminated with Fusarium moniliforme)
- IT 116355-83-0, Fumonisin B1 116355-84-1
RL: BIOL (Biological study)
(of **corn** contaminated with Fusarium moniliforme, **toxicity** of, equine leukoencephalomalacia in relation to)
- IT 9031-66-7, Transaminase
RL: PROC (Process)
(of serum, increase of, by **dietary corn** contaminated with Fusarium moniliforme)
- L71 ANSWER 17 OF 28 HCAPLUS COPYRIGHT 2003 ACS
AN 1989:626957 HCAPLUS
DN 111:226957
TI Production of fusarenone X, nivalenol, and zearalenone by Gibberella zeae isolates, and their **toxicity** in fibroblasts and **rats**
AU Abbas, H. K.; Mirocha, C. J.
CS Dep. Plant Pathol., Univ. Minnesota, St. Paul, MN, 55108, USA
SO Mycotoxin Research (1988), 4(2), 67-74
CODEN: MYREET; ISSN: 0178-7888
DT Journal
LA English
CC 4-5 (Toxicology)
Section cross-reference(s): 17
- AB Three isolates of G. zeae, the perfect stage of Fusarium graminearum, were isolated from ground **corn** cultures obtained from Taiwan in 1985 and identified as G. zeae 1.1, G. zeae 1-5, and G. zeae 1-7. The isolates were grown on a solid rice medium and exts. prepd. with 75% aq. methanol. The exts. were examd. for **toxicity** in the following systems: cytotoxicity to cultured normal human diploid skin fibroblasts and **mouse** fibroblasts, and **toxicity** to **rats** of unextd. cultures. The three exts. were highly cytotoxic as indicated by the ability to cause death and disintegration of 3T3 Swiss **mouse** fibroblasts and human diploid skin fibroblasts during 3 to 4 days in culture. The unextd. cultures of the isolates were highly **toxic** to **rat**, causing hemorrhage of tissues (bladder, **stomach**, and **intestine**), uterine enlargement, small thymuses, small spleens, wt. loss, and death. The exts. were tested for prodn. of trichothecenes (nivalenol and fusarenone X) and zearalenone on rice grains. Prodn. of the 3 mycotoxins was greater at room temp. than in the cold room. Detection of the 3 mycotoxins from the cultures was variable, ranging from 273 to 817 ppm for nivalenol, 268 to 662 ppm for fusarenone, and 162 to 1095 ppm for zearalenone at room temp.,

- and 159 to 413 ppm for nivalenol, 113 to 125 ppm for fusarenone X, and 44 to 202 ppm for zearalenone in the cold room (10.degree.).
- ST fusarenone X Gibberella fibroblast; nivalenol **toxicity** fibroblast; zearalenone **toxicity** fibroblast; mycotoxin Gibberella fibroblast
- IT Gibberella zeae
(mycotoxins prodn. by, **toxicity** of, to fibroblasts)
- IT Skin, **toxic** chemical and physical damage
(mycotoxins **toxicity** to fibroblasts of, of human)
- IT Fibroblast
(mycotoxins **toxicity** to, of humans and lab. animals)
- IT **Toxicity**
(of mycotoxins, to human and lab. animal fibroblasts)
- IT Mycotoxins
RL: PREP (Preparation)
(prodn. of, by Gibberella zeae, human and lab. animal fibroblast **toxicity** in relation to)
- IT 17924-92-4P, Zearalenone 23255-69-8P, Fusarenone X 23282-20-4P, Nivalenol
RL: PREP (Preparation)
(prodn. of, by Gibberella zeae, human and lab. animal fibroblast **toxicity** in relation to)
- L71 ANSWER 18 OF 28 HCAPLUS COPYRIGHT 2003 ACS
- AN 1987:472327 HCAPLUS
- DN 107:72327
- TI Effects of polybrominated biphenyls upon **rat** urinary protein patterns as detected by two-dimensional electrophoresis
- AU Myrick, James E.; Robinson, Mary K.; Hubert, Ivey Lois; Smith, S. Jay; Hannon, W. Harry
- CS Cent. Environ. Health, U.S. Dep. Health Hum. Serv., Atlanta, GA, 30333, USA
- SO Archives of Environmental Contamination and Toxicology (1987), 16(5), 579-85
CODEN: AECTCV; ISSN: 0090-4341
- DT Journal
- LA English
- CC 4-3 (Toxicology)
- AB Five-month-old female **rats** were dosed by gavage with 0 or 500 mg of Fire Master BP 6 (polybrominated biphenyls, PBBs)/kg as a 5% soln. in **corn** oil and then maintained on normal **diet** and water **ad libitum**. Urine pools from **test** and control groups of 5 fasting **rats** were collected once a month, up to 179 days after the **test rats** were dosed. PBB-dosed animals showed significantly higher creatinine-cor. proteinuria. Two-dimensional electrophoresis (2DE) and Ag staining were used to det. differences in urinary protein patterns between **test** and control **rats**. Dosed animals excreted new proteins with higher mol. wt. and more basic isoelec. points than controls. The appearance of numerous, new proteins excreted by the PBB-dosed **rats** demonstrates the utility of 2DE for the sensitive detection of possible health effects from **toxicant** exposure.
- ST brominated biphenyl urine protein; proteinuria polybrominated biphenyl
- IT Urine
(proteins excretion in, after polybrominated biphenyl administration, electrophoresis for study of)
- IT Proteins, biological studies
RL: BIOL (Biological study)

(metabolic disorders, proteinuria, from polybrominated biphenyls, electrophoresis for study of)

IT 92-52-4D, Biphenyl, polybrominated biphenyls
RL: BIOL (Biological study)
(proteinuria from, electrophoresis for study of)

L71 ANSWER 19 OF 28 HCAPLUS COPYRIGHT 2003 ACS
AN 1987:470372 HCAPLUS
DN 107:70372
TI Preliminary **toxicity** profile of arotinoids SMR-2 and SMR-6 in male B6D2F1 **mice**
AU Lindamood, Charles, III; Giles, Herschell D.; Hill, Donald L.
CS Kettering-Meyer Lab., South. Res. Inst., Birmingham, AL, 35255-5305, USA
SO Fundamental and Applied Toxicology (1987), 8(4), 517-30
CODEN: FAATDF; ISSN: 0272-0590
DT Journal
LA English
CC 1-6 (Pharmacology)
Section cross-reference(s): 4
GI



I, R=CH₂OH

II, R=CO₂H

AB Arotinoids, which are analogs of retinoic acid (RA) and and retinol (RO) with the carbon skeleton in a rigid conformation, have more favorable therapeutic indexes relative to all-trans-RA and all-trans-RO. The purpose of this investigation was to obtain preliminary in vivo **toxicity** data on SMR-2 (I; analog of RO) and SMR-6 (II; analog of RA), arotinoids with promising activity for reversal of keratinization in tracheal organ culture. A preliminary **toxicity** study was conducted in male B6D2F1 **mice** with gavage of retinoids in corn oil (0.01, 0.05, and 0.1 mg/kg-day of I or II; 1,5, and 10 mg/kg-day of RA as ref. control). Due to lack of **toxicity**, each dose level for I and II was increased by 4-fold on day 29 of dosing. The study was terminated on day 57. Hypervitaminosis A (**wt. loss**, alopecia, skin scaling, and bone thinning) was induced in the mild- and high-dose SMR groups; **wt.-gain** depression was predominant in the high-dose RA group. The SMR compds. were .apprx.100-fold more **toxic**, based on **wt. loss**, than RA. In the SMR dose groups with hypervitaminosis A, white blood cell counts were elevated 2-4-fold; and there were microscopic lesions in skin, **testes**, epididymis, bone, thymus, bone marrow, peripheral lymph nodes, spleen, **stomach**, adrenal, and pituitary. The leukocytosis was attributed to leukopoiesis in spleen and bone marrow, which may be due to either a direct effect and/or a secondary response to a subacute inflammatory reaction in skin. Only peripheral lymph node hyperplasia was obsd. in I and RA low-dose groups. Enlarged thymus, lymph node hyperplasia, leukopoiesis in spleen and bone marrow, elevated alk. phosphatase with bone hypertrophy, and **testicular** degeneration were obsd. in the mid-dose RA group. The results indicate that immune stimulation may be a

primary early response to retinoids and that skin, leukopoietic tissues, reproductive organs, and bone are primary targets for retinoid toxicity.

ST arotinoid SMR2 SMR6 toxicity

IT Toxicity

(of arotinoids)

IT 89315-18-4 109791-92-6

RL: ADV (Adverse effect, including toxicity); BIOL (Biological study) (toxicity of)

L71 ANSWER 20 OF 28 HCAPLUS COPYRIGHT 2003 ACS

AN 1986:547953 HCAPLUS

DN 105:147953

TI Toxicology and carcinogenesis studies of 3-chloro-2-methylpropene (technical grade containing 5% dimethylvinyl chloride) (CAS No. 563-47-3) in F344/N rats and B6C3F1 mice. (Gavage studies)

CS National Toxicology Program, Research Triangle Park, NC, 27709, USA

SO Natl. Toxicol. Program Tech. Rep. Ser. (1986), 300, 196 pp.

CODEN: NTPSE3

DT Report

LA English

CC 4-6 (Toxicology)

AB Toxicol. and carcinogenesis studies of tech.-grade 3-chloro-2-methylpropene (I) [563-47-3] (contg. 5% dimethylvinyl chloride [513-37-1]), a widely used insecticide and chem. intermediate, were performed on rats and mice. In the 13-wk studies, 50%-100% mortality occurred in groups of male and female rats receiving 400 mg/kg, male rats receiving 300 mg/kg, and male and female mice receiving 500-1250 mg/kg. Inflammation and necrosis of the liver were seen in rats and mice, and necrosis of cortical tubules of the kidney was seen in mice. Based on these observations, groups of 50 male and 50 female rats were administered I in corn oil by gavage at doses of 0, 75, or 150 mg/kg, 5 days/wk for 103 wk, and groups of 50 male and 50 female mice received I at 0, 100, or 200 mg/kg on the same schedule. In the 2-yr studies, the mean body wt. of high-dose male rats was consistently 10-15% lower than that of the vehicle control group, and late in the study there was a marginal redn. in survival of high-dose male rats. Mean body wts. and survival in low-dose male rats and in both dosed groups of female rats were comparable to those of their vehicle control groups. Mean body wts. of high-dose male mice and of both dosed groups of female mice were slightly (5-9%) lower than those of the vehicle controls, whereas survival in both male and female mice was not affected by I administration. Dose-related increases in the incidence of forestomach inflammation were obsd. in male and female mice (male: vehicle control, 0/49; low dose, 9/49; high dose, 7/49; female: vehicle control, 2/50; low dose, 3/48 high dose, 9/44). Increased incidences of forestomach basal cell hyperplasia were obsd. in rats and mice of each sex. I induced forestomach squamous cell papillomas and squamous cell carcinomas in rats and mice. Invasion or metastasis of the squamous cell carcinomas to other organs was obsd. in 2 low-dose male, 3 high-dose male, and 1 high-dose female mice. Renal tubular cell adenocarcinomas (1/49), renal transitional cell carcinomas (1/49), and transitional cell papillomas (1/46) of the urinary bladder were obsd. in high-dose male rats, and renal tubular cell adenomas (1/50) and renal tubular cell

adenocarcinomas (1/50) were seen in low-dose male **rats**. These urinary tract neoplasms were not obsd. in vehicle controls. The incidences of inflammation of the nasal cavity and of nephropathy/nephrosis were greater in the 2 dosed groups than in the vehicle control groups of **rats** and **mice** of each sex. Neg. trends or lower incidences of pheochromocytomas of the adrenal gland and C-cell adenomas or carcinomas (combined) of the thyroid gland were obsd. in dosed male **rats**. Neg. trends or lower incidences of pheochromocytomas of the adrenal gland and C-cell adenomas or carcinomas (combined) of the thyroid gland were obsd. in dosed male **rats**. I was inactive or only weakly mutagenic in expts. with several strains of Salmonella typhimurium, but it was mutagenic in the **mouse** lymphoma L5178Y/TK+/- forward mutation assay without exogenous metabolic activation. Cytogenetics **tests** with cultured Chinese hamster ovary cells were pos. for induction of chromosomal aberrations and sister-chromatid exchanges (SCE's) in the absence of **rat** liver S9. With metabolic activation, SCE levels remained significantly elevated, but the no. of chromosomal aberrations was reduced. Under the conditions of these 2-yr gavage studies, there was clear evidence of carcinogenicity for I as shown by the increased incidences of squamous cell neoplasm in the forstomach of male and female **rats** and **mice**.

- ST chloromethylpropene dimethylvinyl chloride **toxicity**;
carcinogenicity chloromethylpropene dimethylvinyl chloride; mutagenicity
chloromethylpropene dimethylvinyl chloride
- IT Sex
(chloromethylpropene carcinogenicity and mutagenicity and
toxicity in relation to)
- IT Adrenal gland, neoplasm
Carcinoma
Kidney, neoplasm
Liver, neoplasm
Lymphoma
Mutation
Neoplasm
Papilloma
(from chloromethylpropene contg. dimethylvinyl chloride as contaminant)
- IT Body **weight**
(**loss** of, from chloromethylpropene contg. dimethylvinyl
chloride as contaminant)
- IT Carcinoma
(adeno-, from chloromethylpropene contg. dimethylvinyl chloride as
contaminant)
- IT **Stomach**, neoplasm
(fore-, from chloromethylpropene contg. dimethylvinyl chloride as
contaminant)
- IT Bladder
Nose
Organ
(neoplasm, from chloromethylpropene contg. dimethylvinyl chloride as
contaminant)
- IT 513-37-1
RL: BIOL (Biological study)
(carcinogenicity and mutagenicity and **toxicity** of tech. grade
chloromethylpropene contg.)
- IT 563-47-3
RL: ADV (Adverse effect, including toxicity); BIOL (Biological study)
(carcinogenicity and mutagenicity and **toxicity** of, contg.)

dimethylvinyl chloride as contaminant)

- L71 ANSWER 21 OF 28 BIOSIS COPYRIGHT 2003 BIOLOGICAL ABSTRACTS INC.
AN 1986:432943 BIOSIS
DN BA82:99131
TI EFFECTS OF LEARNED FLAVOR AVOIDANCE ON GROOMING BEHAVIOR IN RATS.
AU REIDINGER R F JR; MASON J R
CS MONELL CHEMICAL SENSES CENTER, 3500 MARKET ST., PHILADELPHIA, PA 19104.
SO PHYSIOL BEHAV, (1986) 37 (6), 925-932.
CODEN: PHBHA4. ISSN: 0031-9384.
FS BA; OLD
LA English
AB In Experiment 1, rats were conditioned to avoid saccharin in tapwater by pairing it with LiCl in **carboxymethylcellulose** (CMC) applied to the fur. Conditioned flavor avoidance (CFA) of saccharin was then assessed in drinking and grooming tests. In Experiment 2, rats were given saccharin CMC on their fur and NaCl in water (or vice-versa) as conditioned stimuli in a CFA paradigm. Two-choice tests (saccharin vs. NaCl) followed in drinking and grooming contexts. In Experiment 3, rats were given saccharin CMC on one flank and vehicle (CMC only) on the other. After grooming, animals were injected with LiCl and then given 2-choice test, first between saccharin and water, then between saccharin-CMC and plain-CMC, and finally, between saccharin and water. Strong CFA was exhibited in drinking tests in all 3 experiments. This was not the case in grooming tests. Rats continued to groom when tastant was applied to only one flank (Experiment 1), and exhibited only weak CFA when a different tastant was applied to each flank (Experiments 2 and 3). We conclude that grooming can be directed to minimize the ingestion of noxious substances, but that such ingestion is not sufficiently reduced to affect the efficacy of grooming as a delivery method for unpalatable substances (e.g., **rodenticides**, chemosterilants). We speculate that grooming represents a weakness in rodents' defenses against dietary poisoning, and it might be used to deliver toxicants as part of crop protection schemes that make use of CFA.
- CC Behavioral Biology - Animal Behavior *07003
Behavioral Biology - Conditioning *07005
Nervous System - Physiology and Biochemistry 20504
Psychiatry - Psychophysiology *21003
BC Muridae 86375
IT Miscellaneous Descriptors
CONDITIONING RODENT CONTROL
- L71 ANSWER 22 OF 28 HCAPLUS COPYRIGHT 2003 ACS
AN 1979:469446 HCAPLUS
DN 91:69446
TI Effects of intubating **rats** with fractions from thermally oxidized **corn** oil and olive oil
AU Gabriel, H. G.; Alexander, J. C.; Valli, V. E.
CS Dep. Nutr., Univ. Guelph, Guelph, ON, N1G 2W1, Can.
SO Nutrition Reports International (1979), 19(4), 515-26
CODEN: NURIBL; ISSN: 0029-6635
DT Journal
LA English
CC 4-3 (Toxicology)
Section cross-reference(s): 17
AB Male weanling **rats** were maintained on Purina lab. chow **ad libitum** throughout the exptl. period, and received either 0.5 mL/day of the distillable nonurea adductable material (DNUA) of

thermally oxidized **corn** oil (CO) or olive oil (OO), or the same amt. of the fresh fats by **stomach** tube. Within 3 days, the **rats** receiving DNUA-OO had lost wt. and were moribund. The **rats** intubated with DNUA-CO gained wt. and were indistinguishable in appearance from the control animals. Addnl. urea adductions of DNUA-CO yielded DNUA-2-CO, which was comparable in **toxicity** to DNUA-OO. The animals in both the latter groups had severe diarrhea, seborrhea and flatulence. Pathol. examn. revealed visceral edema, gastric ulcers, and yeast infection of the **gut**. Histopathol. examn. of tissues showed diffuse lesions. Livers and kidneys of the 3 **test** groups were increased in wt. compared to their resp. controls. Extensive changes in constituent fatty acids of the neutral and polar organ lipids were obsd. Overall evaluation of the thermally oxidized samples showed that DNUA-2-CO and DNUA-OO were very **toxic**, whereas DNUA-CO compared favorably with the control fats.

ST **corn** oil thermal oxidn **toxicity**; olive oil thermal oxidn **toxicity**

IT Fats, biological studies

RL: RCT (Reactant); RACT (Reactant or reagent)
(thermal oxidn. of, **toxicity** in relation to)

IT **Corn** oil
Olive oil

RL: BIOL (Biological study)
(thermally-oxidized, **toxicity** of)

L71 ANSWER 23 OF 28 HCAPLUS COPYRIGHT 2003 ACS

AN 1977:512696 HCAPLUS

DN 87:112696

TI Myocardial lesions in **rats** fed rapeseed oil. I. Influence of strain of **rats**

AU Hulan, H. W.; Kramer, J. K. G.; Corner, A. H.

CS Res. Branch, Agric. Canada, Ottawa, ON, Can.

SO Canadian Journal of Physiology and Pharmacology (1977), 55(2), 258-64
CODEN: CJPPA3; ISSN: 0008-4212

DT Journal

LA English

CC 4-6 (Toxicology)

Section cross-reference(s): 17

AB The influence of strain of **rat** on the development of myocardial lesions was investigated in an expt. which included 2 factors: strain (Wistar, Sherman, Chester Beatty (Hooded), and Sprague-Dawley from 2 sources designated Sprague-Dawley A, Sprague-Dawley C) and **diet** (5% **corn** oil, 20% **corn** oil, and 20% Brassica napus var. Zephyr rapeseed oil). Groups of 30 **rats**, housed 2 per cage, from each of 3 different strains of **rats** and groups of 10 **rats**, housed 2 per cage, from one strain of **rats** (Hooded) were fed semisynthetic **diets** contg. the **test** oils for 16 weeks on an **ad libitum** basis. **Rats** of the Hooded strain consumed considerably less **feed** and grew at a significantly slower rate than did **rats** of the Sprague-Dawley C strain, which in turn ate less and grew significantly slower than **rats** of the other 3 strains. No marked differences were obsd. in the fatty acid compn. of total cardiac lipids among strains. **Rats** from all strains except Hooded fed the **diet** contg. 20% Zephyr RSO had a significantly higher incidence of myocardial lesions than **rats** fed the 20% **corn** oil **diet** which in turn had a significantly higher incidence than **rats** fed 5% **corn** oil. Similarly, significantly more Sprague-Dawley C

rats had myocardial lesions than Sprague-Dawley A **rats** regardless of **diet**. None of the Hooded **rats** fed the **diet** contg. 20% Zephyr RSO developed myocardial lesions whereas all other strains fed this **diet** developed a high incidence of myocardial lesions.

ST myocardial lesion rapeseed oil; heart damage rapeseed oil

IT Heart, **toxic** chemical and physical damage
(myocardial lesions from rapeseed oil in, **rat** strains in relation to)

IT Rape oil

RL: BIOL (Biological study)
(myocardial lesions from, **rat** strains in relation to)

L71 ANSWER 24 OF 28 HCAPLUS COPYRIGHT 2003 ACS

AN 1972:139084 HCAPLUS

DN 76:139084

TI Nutritive protein from **cellulose**

IN Srinivasan, Vadake R.; Callihan, Clayton D.

PA Louisiana State University Foundation

SO U.S., 4 pp.

CODEN: USXXAM

DT Patent

LA English

IC C12D

NCL 195033000

CC 16 (Fermentations)

Section cross-reference(s): 17, 18

FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI US 3627095	A	19711214	US 1969-847719	19690805
PRAI US 1969-847719		19690805		

AB Nutritive protein is prepd. from **cellulose** by cultivating a cellulase-elaborating microorganism and *Alcaligenes faecalis* on delignified **cellulose** under submerged aerobic conditions at pH 5-9 and temp. 25-65.degree.. Thus, 760 parts of sugarcane bagasse was slurried in 8% NaOH for 1 hr at room temp. Excess liq. was removed and the residual solid was spread in a thin layer on a wire **screen** and heated in a hot air oven at 110.degree. for 15 min. The partially dried material was dild. to 193.1 l. with a sterile aq. soln. contg. NaCl 900, (NH₄)₂SO₄ 194, KH₂PO₄ 97, K₂HPO₄ 97, MgSO₄ 19.4, CaCl₂ 19.4, and yeast ext. 60 g. Sufficient HCl to neutralize and to keep the soln. at the neutral point was added. The mixt. was boiled 3 hr and air was introduced slowly for cooling. When the temp. reached 32.degree., 1 l. of inoculum contg. *Cellulomonas* species ATCC No. 21399 and *A. faecalis* ATCC No. 21400 was added. With aeration at 34 l./min, fermentation was carried on for 90.25 hr at 32.degree.. After sepg. unreacted **cellulose** and harvesting the bacterial cells produced, 738 g (97.2%) of the initial insol. **cellulose** had been solubilized and a yield of 370 g of bacterial cells was obtained. The cells contained 50.2% protein. The protein, hydrolyzed in 6N HCl for 22 hr at 110.degree., contained arginine 5.3, histidine 2.3, isoleucine 6.0, leucine 10.0, lysine 6.5, methionine 2.0, phenylalanine 5.8, tyrosine 2.7, threonine 8.3, and valine 5.6%. The digestibility of the protein was detd. by **feeding** weanling **rats ad libitum** over a 10-day period with a protein-free **diet** and the same **diet** with 10, 20, or 40% of the produced protein. All the **rats** receiving 20% or more protein in the **diet** showed a wt. increase, while those on the

protein-free diet showed a net wt. loss. No toxicity occurred at any feeding level.

ST protein nutrition cellulose Alcaligenes

IT Cellulomonas
(cultivation of Alcaligenes faecalis and, on cellulose)

IT Alcaligenes faecalis
(cultivation of Cellulomonas and, on cellulose)

IT Bagasse
(fermentation of, protein manuf. by)

IT Proteins
RL: BIOL (Biological study)
(single-cell, from cellulose)

IT 9004-34-6, biological studies
RL: BIOL (Biological study)
(fermentation of, protein manuf. by)

L71 ANSWER 25 OF 28 HCAPLUS COPYRIGHT 2003 ACS

AN 1967:103813 HCAPLUS

DN 66:103813

TI Relation between cerebral sulfhydryl concentration and the central action of some thio compounds

AU Cascio, Giovanni; Palazzoadriano, Mario; Madonia, Paolino

CS Univ. Palermo, Sicily, Italy

SO Biochimica e Biologia Sperimentale (1966), 5(1), 3-9

CODEN: BBSPAJ; ISSN: 0006-2995

DT Journal

LA Italian

CC 15 (Pharmacodynamics)

AB The effect of cysteamine (I), BAL (II), cysteamine-N-acetic acid (III), and cysteine (IV), alone or in assocn. with substances capable of influencing oxidn.-redn. processes (vitamin B12 (V) and dinitrophenol (VI)), on behavior and cerebral and liver sulfhydryl (SH) concn. was studied in rats and doves, maintained under standard conditions and diets, the doves being given maize and water ad libitum. The compds. were injected into the 24-hr. fasted animals, i.p. in the rats and i.m. in the doves. When V and VI were used, they were administered 30 min. before the thio compds. Thirty min. after injection of the thio compds. the animals were killed by decapitation, bled, and the brain and liver removed. Portions of 300 mg. were homogenized in water and the SH detd. iodometrically (Merville, et al., CA 55, 9531i). With I the ED100 was 200 mg./kg. and the min. convulsant level of SH was a 24% increase above normal. Below this dose brain and liver SH levels were not significantly affected, even at 180 mg./kg. At 200 mg./kg. if the rats were killed before onset of convulsions, cerebral SH was not affected, but if killed during convulsions, brain SH markedly increased (33%) while liver SH was not affected. Pretreatment with V or VI (10 and 20 mg./kg. resp.) which alone did not modify the SH levels, produced convulsions and increases in cerebral SH with the normally non-convulsant dose of 180 mg./kg. In doves I produced no convulsions even at 300 mg./kg. and did not affect cerebral free SH with or without V or VI. II gave similar results; 45 mg./kg. had no effect, 80 mg./kg. produced convulsions and a 33% rise in cerebral and liver SH in rats. With 10 mg./kg. V, 45 mg./kg. produced the same effect as 80 mg./kg. used alone. III even in toxic doses showed a depressive action in all species tested and in rats at 800 mg./kg. did not appreciably affect cerebral SH but increased liver SH distinctly. Pretreatment with V or VI did not modify the effects. IV at 500 mg./kg. produced no appreciable effect but 1000

mg./kg. produced depressive symptoms with 16% rise in cerebral SH. The symptoms lasted 2-3 hrs., then cleared, leaving the animal apparently normal. Pretreatment with V had no effect but that with VI provoked excitatory phenomena with 52% rise in cerebral SH. It is presumed that the convulsant capacity depends on the ratio of velocity of passage through the blood-brain barrier and the rate of oxidn. of nervous tissue and that V and VI accentuate the convulsant power by favoring the permanence of a state of redn. and permitting the concn. of SH in the central nervous system to reach concns. sufficiently higher than normal. 11 references.

ST SULFHYDRYLS BRAIN; BRAIN SULFHYDRYLS; LIVER SULFHYDRYLS; THIO COMPDs
TISSUE SULFHYDRYLS; CONVULSANTS THIO COMPDs; VITAMIN B12 CONVULSIONS

IT Mercapto group

(in brain after thio compd. administration, behavior and)

IT Brain, composition

Liver, composition

(mercapto group in, thio compd. effect on)

IT Behavior

(thio compd. effect on, mercapto groups in brain and)

IT 51-28-5, biological studies

RL: BIOL (Biological study)

(behavior and mercapto group in brain after treatment with, thio compds. and)

IT 52-90-4, biological studies 59-52-9 60-23-1 3724-83-2

RL: BIOL (Biological study)

(behavior and mercapto groups in brain after treatment with)

IT 68-19-9, biological studies

RL: BIOL (Biological study)

(behavior in mercapto groups in brain after treatment with, thio compds. and)

L71 ANSWER 26 OF 28 HCAPLUS COPYRIGHT 2003 ACS

AN 1964:48411 HCAPLUS

DN 60:48411

OREF 60:8542e-h

TI **Toxicological** and hygienic characterization of new chlorinated insecticides and food products contaminated by them

AU Serebryanaya, S. G.

SO Gigiena i Toksikol. Novykh Pestitsidov i Klinika Otravlenii, Dokl. 2-oi [Vtoroi] Vses. Konf. (1962) 124-33

DT Journal

LA Unavailable

CC 69 (Toxicology, Air Pollution, and Industrial Hygiene)

AB The effect of dichlorodiphenyldichloroethane (I), **diethyldiphenyldichloroethane** (II), dimethoxydiphenyltrichloroethane (III), and dichlorodiphenyltrichloroethane (IV) on the behavior of animals, body wt., duration of intoxication, reproduction and succeeding generations, morphological compn. of blood and changes in internal organs, phys.-chem. changes in urine, glycemia, carbohydrate metabolism in liver, Thunberg's test, and changes of relative wt. of organs was examd. in **rats**, **mice**, and cats. The minimal **toxic** dose, and L.D.min. of I, II, III, and IV was 1000, 2000, 3000, and 150 mg./kg. and 3000, >4000, >5000, and 300 mg./kg., resp., for **rats**; -, 1000, -, and 150 mg./kg. and 3000, >4000, 4000, and 200 mg./kg., resp., for cats; and 2000, 2000, 2000, and 150 mg./kg. and 4000, >4000, -, and 200 mg./kg. for **mice**. I, II, and III had no cumulative effect. During 11 months of daily administration of I, II, III, and IV, the minimal physiol. active doses were 10, 25, 10, and 0.2

mg./kg. for rats and 5, 10, 10, and 0.1 mg./kg. for cats. I, II, and III, when administered daily to pregnant animals or during lactation, had no effect on the 2nd generation. IV was **toxic** to lactating animals. After chronic I, II, III, and IV intoxication, anorexia, **loss of wt.**, hypohemoglobinemia, leukocytosis, and tremor were observed. The glycemic curves had a diabetic character only after III or IV application. IV caused a decrease of dehydrogenation activity of the liver. I and III gave a similar, but smaller effect, while II potentiated this function. IV caused an increase of the relative wt. of liver, while I, II, and III did not. The max. permissible amts. in food products were established as 7 mg./kg. in fruit and 3.5 mg./kg. in **corn** for I, 14 and 7 mg./kg. for II, and 14 mg./kg. for all products for III.

- IT Blood
- Nervous system
- (chlorinated insecticide effect on)
- IT Insecticides
- (chlorinated, **toxicology** of)
- IT Liver
- (effect of chlorinated insecticides on)
- IT Food
- (insecticide residues in)
- IT Field theories
- (symmetry, boson decay probabilities and)
- IT Ethane, 1,1,1-trichloro-2,2-bis(p-methoxybenzyl)-
- (**toxicology** of)
- IT 50-99-7, Blood sugar
- (chlorinated-insecticide effect on)
- IT 72-54-8, Ethane, 1,1-dichloro-2,2-bis(p-chlorophenyl)-
- (**toxicity** to *Amblyseius hibisci* and other phytoseiid mites,
- toxicology** of)
- IT 50-29-3, Ethane, 1,1,1-trichloro-2,2-bis(p-chlorophenyl)- 72-56-0,
- Ethane, 1,1-dichloro-2,2-bis(p-ethylphenyl)-
- (**toxicology** of)

L71 ANSWER 27 OF 28 HCAPLUS COPYRIGHT 2003 ACS

AN 1958:98396 HCAPLUS

DN 52:98396

OREF 52:17371e-i,17372a-b

TI Decumbin, a new compound from a species of *Penicillium*

AU Singleton, V. I.; Bohonos, N.; Ullstrup, A. J.

CS Purdue Univ., Lafayette, IN

SO Nature (1958), 181, 1072-3

DT Journal

LA Unavailable

CC 11C (Biological Chemistry: Microbiology)

AB *Penicillium decumbens* isolated during a microbiol. study of **corn** spoilage in storage produced cryst. bodies which when purified and isolated resulted in colorless, odorless needles melting at 204.degree.. The compd. is considered to be a previously unreported dihydroxy, .alpha.-.beta.-unsatd. lactone and has been called decumbin. Production of visible crystals was best on a medium of potato-dextrose broth prepd. by steaming diced, peeled potatoes in 500 ml. tap H2O for 30 min., draining through one layer of cheese cloth, adding this to 20 g. com. glucose and dilg. to 1 l. Stationary surface culture was used since decumbin produced in submerged culture was more difficult to purify. The sterile potato-dextrose broth, in 300 ml. portions per rectangular 2 qt. milk bottle, was inoculated with a heavy spore suspension of *P. decumbens*.

In about 6 days at 23-26.degree., growth appeared complete with uniform grey-green sporulation and orange droplets of fluid on the surface of the mycelium. Needles of decumbin up to 1/4 in. long attached to the submerged side of the mycelium began to appear after 8 days. Cultures were usually harvested 10-12 days. One hundred bottles were strained through two layers of cheese cloth, and the mycelium extd. with 3 l-1. portions of boiling MeOH. The ext. was chilled to 10.degree., the ppt. extd. with 200-300 ml. hot MeOH. The combined exts. were chilled overnight, filtered, dild. with 100 ml. distd. H2O and concd. in vacuo until free of MeOH. The crystals of decumbin were filtered from the aq. soln. Average yield of crude decumbin, m.p. 190.degree. was 278 mg./l. of medium in a total of 1,404 bottles processed. The crystals were purified by washing with petroleum ether (3 ml./gm.) and recrystd. from 50% (v/v) aq. MeOH (33 ml./g.) and ethyl acetate (60 ml./g.). The pure decumbin produced by several recrystns. from these two solvents was not altered by careful sublimation in vacuo or drying in vacuo over phosphoric anhydr. at 100.degree.. The solubility of decumbin at 26.degree., detd. by wt. loss from a single crystal after 19 hr. with frequent stirring, was 0.6. mg./l. of H2O, 1.2 mg./l. of 9.5% EtOH and 24.9 g./l. of absolute EtOH. Preliminary tests showed decumbin to be **toxic** to **rats** and goldfish and inhibitory to wheat germination, but not an active antibiotic under test conditions. It gave no zones of inhibition when 1 mg./l. solns. in 40% (v/v) MeOH in H2O were used in agar cup diffusion tests against Staphylococcus aureus, Escherichia coli, Hormodendrum cladosporoides. The acute oral L.D.50 for **rats** was of the order of 275 mg./kg. Autopsy of **rats** showed congestion of the auricles and assocd. blood vessels with dark blood and fullness of the **stomach** with undigested food. Hydrogenated decumbin was not **toxic** to two **rats** at 2,700 mg./kg. This physiol. activity or **toxicity** of decumbin in mammals, cold-blooded animals, and plants is similar to that found with other unsatd. lactones and the loss of **toxicity** upon hydrogenation is also typical of these compds.

- IT Wheat
(decumben **toxicity** to)
- IT Penicillium decumbens
(dihydroxy .alpha.,.beta.-unsatd. lactone in)
- IT Lactones
(dihydroxy, .alpha.,.beta.-unsatd., from Penicillium decumbens)
- IT 20350-15-6, Decumbin
(from Penicillium decumbens)

L71 ANSWER 28 OF 28 HCAPLUS COPYRIGHT 2003 ACS

AN 1955:70774 HCAPLUS

DN 49:70774

OREF 49:13505b-d

TI Influence of **diet** on the acute **toxicity** of chlorpromazine in **mice**

AU Courvoisier, Simone; Cosar, Charles

SO Compt. rend. (1955), 240, 2026-7

DT Journal

LA Unavailable

CC 11H (Biological Chemistry: Pharmacology)

AB **Mice** were maintained on a balanced synthetic **diet** (I) or on an empirical **diet** (II) in order to test the effect of **diet** on the 50% lethal doses of chlorpromazine (2-chloro-10-(3-dimethylaminopropyl)phenothiazine-HCl) after subcutaneous administration. The L.D.50 on **diet** I for **mice** between

the ages of 90 and 100 days and with a mean wt. of 19.9 g. was 0.43 g./kg.; for **mice** in the same age group with a mean wt. of 24.1 g. on **diet II** (bread, whole milk, **corn** and lettuce, **ad libitum**) the L.D.50 was 0.175 g./kg. For **mice** between the ages of 53 and 55 days with a mean wt. of 16.5 g. the L.D.50 on **diet I** was above 0.35 g./kg.; for **mice** in the same age group with a mean wt. of 20.4 g. the L.D.50 was 0.16 g./kg. on **diet II**. For **mice** between the ages of 50 and 56 days with a mean wt. of 22 g., the L.D.50 on **diet I** was 0.4 g./kg.; for **mice** in the same age group with a mean wt. of 15 g. the L.D.50 was 0.2 g./kg. on **diet II**. For **mice** between the ages of 35 and 40 days with a mean wt. of 18.7 g., the L.D.50 on **diet I** was 0.465 g./kg.; for **mice** in the same age group with a mean wt. of 18.9 g. the L.D.50 was 0.18 g./kg. on **diet II**. These results indicate that regardless of the age or wt. of the **mice** the **toxicity** of chlorpromazine is at least twice as great on an empirical **diet** as on a balanced **diet**

IT **Diet**(chlorpromazine **toxicity** and)IT 50-53-3, Phenothiazine, 2-chloro-10-(3-dimethylaminopropyl)-
(therapy with)IT 50-53-3, Phenothiazine, 2-chloro-10-(3-dimethylaminopropyl)-
(**toxicity** of, **diet** and)